arXiv:1105.0160v2 [math.QA] 5 May 2011

# Classification of Finite Dimensional Irreducible Representations of Generalized Quantum Groups via Weyl Groupoids

Saeid Azam, Hiroyuki Yamane, Malihe Yousofzadeh

### Abstract

Let $U(\chi)$ be a generalized quantum group such that $\dim U^+(\chi) = \infty$, $|R^+(\chi)| < \infty$, and $R^+(\chi)$ is irreducible, where $U^+(\chi)$ is the positive part of $U(\chi)$, and $R^+(\chi)$ is the Kharchenko's positive root system of $U^+(\chi)$. In this paper, we give a list of finite-dimensional irreducible highest weight $U(\chi)$-modules, relying on a *special* reduced expression of the longest element of the Weyl groupoid of $U(\chi)$.

## Introduction

In this paper, we give a list of the finite-dimensional irreducible highest weight modules of a generalized quantum group $U(\chi)$ whose positive part $U^+(\chi)$ is infinite-dimensional and has a Kharchenko's PBW-basis with a finite irreducible positive root system. We call such $U(\chi)$ *finite-type infinite-dimensional* generalized quantum group.

We begin with recalling some facts of Lie superalgebras. The *contragredient Lie superalgebras* [11, Subsection 2.5.1] is defined in a way similar to that for Kac-Moody Lie algebras. Kac classified the simple Lie superalgebras [11, Theorem 5], where the finite-dimensional irreducible contragredient Lie superalgebras played crucial roles; those are

(1) the simple Lie algebras of type $X_N$, where $X = A, \ldots, G$,
(2) $sl(m+1|n+1)$ $(m+n \geq 2)$,
(3) $B(m,n)$ $(m \geq 0, n \geq 1)$, $C(n)$ $(n \geq 3)$, $D(m,n)$ $(m \geq 2, n \geq 1)$, $D(2,1;x)$ $(x \neq 0, -1)$, $F(4)$, $G(3)$.

The ones in (1) and (3) are simple. The simple Lie superalgebras $A(m,n)$ is defined by $sl(m+1|n+1)$ if $m \neq n$, and otherwise $A(n,n) := sl(n+1|n+1)/\mathfrak{i}$, where $\mathfrak{i}$ is a unique one-dimensional ideal.

Bases of the root system of the Lie superalgebras of (2)-(3) are not conjugate under the action of their Weyl groups. However each two of them are transformed to each other under the action of their *Weyl groupoids* $W$, whose axiomatic treatment was introduced by Heckenberger and the second author [9]. Kac [11, Theorem 8 (c)] gave a list of the finite-dimensional irreducible highest weight modules of the simple Lie superalgebras in (2)-(3) above. In the same way as that for the proof of our main result, Theorem 18, we can have a new proof of recovering the list; our idea is to use a specially good one among the reduced expressions of the longest elements of the Weyl groupoids $W$.

Let $\mathfrak{g} := sl(m+1|n+1)$ or $C(n)$ for example. Let $\mathfrak{h}$ be a Cartan subalgebra of $\mathfrak{g}$ such that the Dynkin diagram of $(\mathfrak{g}, \mathfrak{h})$ is a standard one. Let $\Pi = \{\alpha_i | 1 \leq i \leq \dim \mathfrak{h}\}$ be the set of simple roots $\alpha_i$ corresponding to $\mathfrak{h}$. Let $w_0$ be the longest element of $W$ of $\mathfrak{g}$ whose end domain is corresponding to $\mathfrak{h}$. The key fact used in this paper is that there exists a reduced expression $s_{i_1} \cdots s_{i_{\ell(w_0)}}$ of $w_0$ such that $s_{i_1} \cdots s_{i_{x-1}}(\alpha_{i_x})$, $1 \leq x \leq k$, are even roots, and $s_{i_1} \cdots s_{i_{y-1}}(\alpha_{i_y})$, $k+1 \leq y \leq \ell(w_0)$, are odd roots for some $k$. We claim that this is essential to the fact that an irreducible highest weight $\mathfrak{g}$-module of a highest weight $\Lambda$ is finite-dimensional if and only if $\frac{2\langle \Lambda, \alpha_i \rangle}{\langle \alpha_i, \alpha_i \rangle} \in \mathbb{Z}_{\geq 0}$ for all even simple roots $\alpha_i$, where $\langle \, , \rangle$ is a bilinear form coming from the Killing form of $\mathfrak{g}$.

Motivated by Andruskiewitsch and Schneider's theory [3], [4] toward classification of pointed Hopf algebras, Heckenberger [7] classified the Nichols algebras of diagonal-type. Let $U(\chi)$ be the $\mathbb{C}$-algebra (we call it the generalized quantum group) defined in the same manner as in the Lusztig's book [13, 3.1.1 (a)-(e)] for any bi-homomorphism $\chi : \mathbb{Z}\Pi \times \mathbb{Z}\Pi \to \mathbb{C}^\times$. We say that $U(\chi)$ is finite-type if the Kharchencko's positive root system of $U(\chi)$ is finite and irreducible. Nichols algebra of diagonal-type is isomorphic to the positive part $U^+(\chi)$ of $U(\chi)$ of finite type. If $U(\chi)$ is finite-type infinite-dimensional, then it is a multi-parameter quantum algebra of a simple Lie algebra in (1), a multi-parameter quantum superalgebra of a simple Lie superalgebra in (2) or (3), or one of two algebras [7, Table 1-Row 5, Table 3-Row 14]. Our main result, Theorem 18, gives a list of the finite dimensional irreducible highest weight modules of such $U(\chi)$ in the way mentioned as above.

Studying representation theory of $U(\chi)$ must be interesting and fruitful since the factorization formula of Shapovalov determinants of any finite-type $U(\chi)$ has been obtained by Heckenberger and the second author in [10]. We believe that it help give us a new way to study Lusztig's conjecture.

This paper is organized as follows. In Section 1, we give a definition of generalized quantum groups $U(\chi)$ associated with any bi-homomorphism $\chi$, and explain the Kharchenko's PBW-theorem of $U(\chi)$. In Section 2, we explain properties of Weyl groupoids associated with finite-type $U(\chi)$, and Heckenberger's classification of finite-type infinite-dimensional $U(\chi)$. Assume that $U(\chi)$ is finite-type infinite-dimensional. In Section 3, using longest elements of Weyl groupoids and

Heckenberger's Lusztig isomorphisms, we give a criterion when an irreducible highest $U(\chi)$-module is finite-dimensional. In Section 4, we give a list of finite-dimensional irreducible highest weight $U(\chi)$-modules for $U(\chi)$ having a standard Dynkin diagram.

In [19], we have given a result similar to Theorem 18 for [7, Table 1-Row 5].

# 1 Generalized quantum groups

## 1.1 Basic terminology

Let $\mathbb{N}$ denote the set of positive integers. Let $\mathbb{Z}$ denote the ring of integers. For $m$, $n \in \mathbb{Z}$, let $J_{m,n} := \{i \in \mathbb{Z} | m \leq i \leq n\}$. For $n \in \mathbb{Z}$, let $J_{n,\infty} := \{i \in \mathbb{Z} | i \geq n\}$, and let $J_{-\infty,n} := \{i \in \mathbb{Z} | i \leq n\}$. Then $\mathbb{N} = J_{1,\infty}$. Let $\mathbb{Z}_{\geq 0} := J_{0,\infty}$. Let $\mathbb{C}$ denote the field of complex numbers. Let $\mathbb{C}^\times := \mathbb{C} \setminus \{0\}$.

For a unital $\mathbb{C}$-algebra $\mathcal{G}$, let $\mathrm{Ch}(\mathcal{G})$ denote the set of $\mathbb{C}$-algebra homomorphisms from $\mathcal{G}$ to $\mathbb{C}$.

For $t \in \mathbb{C}^\times$ and $m$, $n \in \mathbb{Z}_{\geq 0}$ with $n \leq m$, let

$$(1.1) \quad (m)_t := \sum_{j \in J_{0,m-1}} t^j, \quad (m)_t! := \prod_{j \in J_{1,m}} (j)_t, \quad \binom{m}{n}_t := \lim_{t' \to t} \frac{(m)_{t'}!}{(n)_{t'}!(m-n)_{t'}!}.$$

For $m \in \mathbb{N}$, let $P_m := \{r \in \mathbb{C}^\times | r^m = 1, r^t \neq 1 (t \in J_{1,m-1})\}$. Let $P_\infty := \mathbb{C}^\times \setminus \cup_{m \in \mathbb{N}} P_m$.

For a $\mathbb{C}$-algebra $\mathfrak{a}$ and $X$, $Y \in \mathfrak{a}$, let $[X, Y] := XY - YX$.

For a set $\mathfrak{s}$, let $|\mathfrak{s}|$ denote the cardinality of $\mathfrak{s}$, and denote by $\mathfrak{S}(\mathfrak{s})$ the set of bijective maps from $\mathfrak{s}$ to $\mathfrak{s}$. Let $\uplus$ mean disjoint union of sets.

For $\mathbb{Z}$-modules $\mathfrak{b}$ and $\mathfrak{c}$, let $\mathrm{Hom}_{\mathbb{Z}}(\mathfrak{b}, \mathfrak{c})$ be the $\mathbb{Z}$-module formed by the $\mathbb{Z}$-module homomorphisms from $\mathfrak{b}$ to $\mathfrak{c}$, and let $\mathrm{Aut}_{\mathbb{Z}}(\mathfrak{b})$ be the group formed by the $\mathbb{Z}$-automorphisms of $\mathfrak{b}$.

The symbol $\delta_{ij}$, or $\delta_{i,j}$, denotes Kronecker's delta, that is, $\delta_{ij} = 1$ if $i = j$, and $\delta_{ij} = 0$ otherwise.

## 1.2 Bi-homomorphism $\chi$ and Dynkin diagram of $\chi$

Throughout this paper, we

(1.2) fix $N \in \mathbb{N}$, let $I := J_{1,N}$, and fix a set $\Pi := \{\alpha_i | i \in I\}$ with $|\Pi| = N$.

Let $\mathbb{Z}\Pi$ be the free $\mathbb{Z}$-module with the basis $\Pi$. Then $\mathrm{rank}_{\mathbb{Z}} \mathbb{Z}\Pi = N$. We say that a map $\chi : \mathbb{Z}\Pi \times \mathbb{Z}\Pi \to \mathbb{C}^\times$ is a *bi-homomorphism on* $\Pi$ if

$$(1.3) \qquad \chi(\alpha, \beta + \gamma) = \chi(\alpha, \beta)\chi(\alpha, \gamma), \quad \chi(\alpha + \beta, \gamma) = \chi(\alpha, \gamma)\chi(\beta, \gamma)$$

hold for all $\alpha$, $\beta$, $\gamma \in \mathbb{Z}\Pi$. Let $\mathcal{X}_\Pi$ be the set of bi-homomorphisms on $\Pi$.

Let $\chi \in \mathcal{X}_\Pi$ and let $q_{ij} := \chi(\alpha_i, \alpha_j)$ for $i, j \in I$. By the *Dynkin diagram* of $\chi$, me mean the un-oriented graph with $N$-dots such that each $i$-th dot is labeled $\alpha_i$ and $q_{ii}$, each two $j$-th and $k$-th dots with $j \neq k$ and $q_{jk}q_{kj} \neq 1$ are joined by a single line labeled $q_{jk}q_{kj}$. For example, if $N = 3$ and $q_{11} = -1$, $q_{22} = \hat{q}^2$, $q_{33} = \hat{q}^6$, $q_{12}q_{21} = \hat{q}^{-2}$, $q_{23}q_{32} = \hat{q}^{-6}$ and $q_{13}q_{31} = 1$ for some $\hat{q} \in P_\infty$, then the Dynkin diagram of $\chi$ is given by the leftmost one of Figure 7.

## 1.3 Quantum group $U = U(\chi)$ associated with $\chi \in \mathcal{X}_\Pi$

From now on until the end of Subsection 1.5, we fix $\chi \in \mathcal{X}_\Pi$, let $q_{ij} := \chi(\alpha_i, \alpha_j)$ for $i, j \in I$.

Let $\tilde{U} = \tilde{U}(\chi)$ be the unital associative $\mathbb{C}$-algebra defined by generators

$$\tilde{K}_\alpha,\, \tilde{L}_\alpha\, (\alpha \in \mathbb{Z}\Pi), \quad \tilde{E}_i, \tilde{F}_i\, (i \in I) \tag{1.4}$$

and relations

$$\begin{array}{l} \tilde{K}_0 = \tilde{L}_0 = 1,\, \tilde{K}_\alpha\tilde{K}_\beta = \tilde{K}_{\alpha+\beta},\, \tilde{L}_\alpha\tilde{L}_\beta = \tilde{L}_{\alpha+\beta},\, \tilde{K}_\alpha\tilde{L}_\beta = \tilde{L}_\beta\tilde{K}_\alpha, \\ \tilde{K}_\alpha\tilde{E}_i = \chi(\alpha, \alpha_i)\tilde{E}_i\tilde{K}_\alpha,\, \tilde{L}_\alpha\tilde{E}_i = \chi(-\alpha_i, \alpha)\tilde{E}_i\tilde{L}_\alpha, \\ \tilde{K}_\alpha\tilde{F}_i = \chi(\alpha, -\alpha_i)\tilde{F}_i\tilde{K}_\alpha,\, \tilde{L}_\alpha\tilde{F}_i = \chi(\alpha_i, \alpha)\tilde{F}_i\tilde{L}_\alpha, \\ \tilde{E}_i\tilde{F}_j - \tilde{F}_j\tilde{E}_i = \delta_{ij}(-\tilde{K}_{\alpha_i} + \tilde{L}_{\alpha_i}) \end{array} \tag{1.5}$$

for all $\alpha, \beta \in \mathbb{Z}\Pi$ and all $i \in I$.

Define the $\mathbb{C}$-algebra automorphism $\tilde{\Omega} : \tilde{U} \to \tilde{U}$ by $\tilde{\Omega}(\tilde{K}_\alpha) := \tilde{K}_{-\alpha}$, $\tilde{\Omega}(\tilde{L}_\alpha) := \tilde{L}_{-\alpha}$, $\tilde{\Omega}(\tilde{E}_i) := \tilde{F}_i\tilde{L}_{-\alpha_i}$, and $\tilde{\Omega}(\tilde{F}_i) := \tilde{K}_{-\alpha_i}\tilde{E}_i$.

Let $\tilde{U}^0 = \tilde{U}^0(\chi)$ be the unital subalgebra of $\tilde{U}$ generated by $\tilde{K}_\alpha$, $\tilde{L}_\alpha$ ($\alpha \in \mathbb{Z}\Pi$). Let $\tilde{U}^+ = \tilde{U}^+(\chi)$ be the unital subalgebra of $\tilde{U}$ generated by $\tilde{E}_i$ ($i \in I$). Let $\tilde{U}^- = \tilde{U}^-(\chi)$ be the unital subalgebra of $\tilde{U}$ generated by $\tilde{F}_i$ ($i \in I$).

**Lemma 1.** *The elements*

$$\tilde{F}_{j_1} \cdots \tilde{F}_{j_m}\tilde{K}_\alpha\tilde{L}_\beta\tilde{E}_{i_1} \cdots \tilde{E}_{i_r} \tag{1.6}$$

*with $m, r \in \mathbb{Z}_{\geq 0}$, $i_x \in I$ ($x \in J_{1,r}$), $j_y \in I$ ($y \in J_{1,m}$), $\alpha, \beta \in \mathbb{Z}\Pi$ form a $\mathbb{C}$-basis of $\tilde{U}$, where we use the convention that if $r = 0$ (resp. $m = 0$), then $\tilde{E}_{i_1} \cdots \tilde{E}_{i_r}$ (resp. $\tilde{F}_{j_1} \cdots \tilde{F}_{j_m}$) means 1.*

*Proof.* This can be proved in a standard way as in [15, Lemma 2.2]. □

Define the $\mathbb{Z}\Pi$-grading structure $\tilde{U} = \oplus_{\alpha\in\mathbb{Z}\Pi}\tilde{U}_\alpha$ on $\tilde{U}$ by $\tilde{K}_\alpha \in \tilde{U}_0$, $\tilde{L}_\alpha \in \tilde{U}_0$, $\tilde{E}_i \in \tilde{U}_{\alpha_i}$, $\tilde{F}_i \in \tilde{U}_{-\alpha_i}$, and $\tilde{U}_\alpha\tilde{U}_\beta \subset \tilde{U}_{\alpha+\beta}$. For $\alpha \in \mathbb{Z}\Pi$, let $\tilde{U}^\pm_\alpha := \tilde{U}^\pm \cap \tilde{U}_\alpha$. Then $\tilde{U}^\pm = \oplus_{\alpha\in\pm\mathbb{Z}_{\geq 0}\Pi}\tilde{U}^\pm_\alpha$.

For $m \in \mathbb{Z}_{\geq 0}$, and $t_1, t_2 \in \mathbb{C}^\times$, let

$$(m; t_1, t_2) := 1 - t_1^{m-1}t_2, \quad (m; t_1, t_2)! := \prod_{j\in J_{1,m}} (j; t_1, t_2). \tag{1.7}$$

For $m \in \mathbb{Z}_{\geq 0}$ and $i$, $j \in I$ with $i \neq j$, define $\tilde{E}^{+}_{m,\alpha_i,\alpha_j}$, $\tilde{E}^{-}_{m,\alpha_i,\alpha_j} \in \tilde{U}^{+}_{m\alpha_i+\alpha_j}$ and $\tilde{F}^{+}_{m,\alpha_i,\alpha_j}$, $\tilde{F}^{-}_{m,\alpha_i,\alpha_j} \in \tilde{U}^{-}_{-m\alpha_i-\alpha_j}$ inductively by $\tilde{E}^{\pm}_{0,\alpha_i,\alpha_j} := \tilde{E}_j$, $\tilde{F}^{\pm}_{0,\alpha_i,\alpha_j} := \tilde{F}_j$, and

$$\begin{aligned}
\tilde{E}^{+}_{m+1,\alpha_i,\alpha_j} &:= \tilde{E}_i\tilde{E}^{+}_{m,\alpha_i,\alpha_j} - q_{ii}^{m}q_{ij}\tilde{E}^{+}_{m,\alpha_i,\alpha_j}\tilde{E}_i, \\
\tilde{E}^{-}_{m+1,\alpha_i,\alpha_j} &:= \tilde{E}_i\tilde{E}^{-}_{m,\alpha_i,\alpha_j} - q_{ii}^{-m}q_{ji}^{-1}\tilde{E}^{-}_{m,\alpha_i,\alpha_j}\tilde{E}_i, \\
\tilde{F}^{+}_{m+1,\alpha_i,\alpha_j} &:= \tilde{F}_i\tilde{F}^{+}_{m,\alpha_i,\alpha_j} - q_{ii}^{m}q_{ji}\tilde{F}^{+}_{m,\alpha_i,\alpha_j}\tilde{F}_i, \\
\tilde{F}^{-}_{m+1,\alpha_i,\alpha_j} &:= \tilde{F}_i\tilde{F}^{-}_{m,\alpha_i,\alpha_j} - q_{ii}^{-m}q_{ij}^{-1}\tilde{F}^{-}_{m,\alpha_i,\alpha_j}\tilde{F}_i.
\end{aligned} \tag{1.8}$$

**Lemma 2.** (1) *For $m \in \mathbb{N}$ and $i$, $j \in I$ with $i \neq j$, we have*

$$[\tilde{E}_i, \tilde{F}_i^{m}] = (m)_{q_{ii}}(-\tilde{K}_{\alpha_i} + q_{ii}^{-m+1}\tilde{L}_{\alpha_i})\tilde{F}_i^{m-1}. \tag{1.9}$$

(2) *For $m \in \mathbb{N}$ and $i$, $j \in I$ with $i \neq j$, we have*

$$[\tilde{E}_i, \tilde{F}^{+}_{m,\alpha_i,\alpha_j}] = -(m)_{q_{ii}}(m; q_{ii}, q_{ij}q_{ji})\tilde{K}_{\alpha_i}\tilde{F}^{+}_{m-1,\alpha_i,\alpha_j}. \tag{1.10}$$

(3) *Let $n$, $m \in \mathbb{Z}_{\geq 0}$ with $n < m$ and $i$, $j \in I$ with $i \neq j$. Then we have*

$$[\tilde{E}^{+}_{n,\alpha_i,\alpha_j}, \tilde{F}^{+}_{m,\alpha_i,\alpha_j}] = (n)_{q_{ii}}!\binom{m}{n}_{q_{ii}}(m; q_{ii}, q_{ij}q_{ji})!\tilde{F}_i^{m-n}\tilde{L}_{n\alpha_i+\alpha_j}. \tag{1.11}$$

*In particular, we have*

$$[\tilde{E}_j^{+}, \tilde{F}^{+}_{m,\alpha_i,\alpha_j}] = (m; q_{ii}, q_{ij}q_{ji})!\tilde{F}_i^{m}\tilde{L}_{\alpha_j}. \tag{1.12}$$

(4) *For $m \in \mathbb{Z}_{\geq 0}$ and $i$, $j \in I$ with $i \neq j$, we have*

$$[\tilde{E}^{+}_{m,\alpha_i,\alpha_j}, \tilde{F}^{+}_{m,\alpha_i,\alpha_j}] = (m)_{q_{ii}}!(m; q_{ii}, q_{ij}q_{ji})!(-\tilde{K}_{m\alpha_i+\alpha_j} + \tilde{L}_{m\alpha_i+\alpha_j}). \tag{1.13}$$

(5) *For $m$, $n \in \mathbb{Z}_{\geq 0}$ and $i$, $j$, $k \in I$ with $i \neq j \neq k \neq i$, we have*

$$[\tilde{E}^{+}_{m,\alpha_i,\alpha_j}, \tilde{F}^{+}_{n,\alpha_i,\alpha_k}] = 0. \tag{1.14}$$

*Proof.* These equations are obtained in a direct way as in [8]. □

For $\alpha = \sum_{i \in I} n_i\alpha_i \in \mathbb{Z}\Pi$ with $n_i \in \mathbb{Z}$, in a way using an induction on $\sum_{i \in I} n_i$, we define the $\mathbb{C}$-subspace $\tilde{\mathcal{I}}^{-}_{-\alpha}$ of $\tilde{U}^{-}_{-\alpha}$ by

$$\tilde{\mathcal{I}}^{-}_{-\alpha} := \begin{cases} \{0\} & \text{if } \alpha \in \{0\} \cup \Pi \cup (\mathbb{Z}\Pi \setminus \mathbb{Z}_{\geq 0}\Pi), \\ \{\tilde{Y} \in \tilde{U}^{-}_{-\alpha} \,|\, \forall i \in I, \, [\tilde{E}_i, \tilde{Y}] \subset \tilde{\mathcal{I}}^{-}_{-\alpha+\alpha_i}\tilde{K}_{\alpha_i} + \tilde{\mathcal{I}}^{-}_{-\alpha+\alpha_i}\tilde{L}_{\alpha_i}\} & \text{otherwise.} \end{cases} \tag{1.15}$$

Let $\tilde{\mathcal{I}}^{-} := \oplus_{\alpha \in \mathbb{Z}_{\geq 0}\Pi \setminus (\{0\} \cup \Pi)}\tilde{\mathcal{I}}^{-}_{-\alpha}$. Let $\tilde{\mathcal{J}}^{-} := \mathrm{Span}_{\mathbb{C}}(\tilde{\mathcal{I}}^{-}\tilde{U}^{0}\tilde{U}^{+})$. Then $\tilde{\mathcal{J}}^{-}$ is an ideal of $\tilde{U}$. We define the unital $\mathbb{C}$-algebra $U = U(\chi)$ by

$$U := \tilde{U}/(\tilde{\mathcal{J}}^{-} + \tilde{\Omega}(\tilde{\mathcal{J}}^{-})) \tag{1.16}$$

(the quotient algebra).

*Remark* 1. (1) By [10, Proposition 3.5 and (3.19)], we can see that the definition of $U(\chi)$ given by (1.16) is equivalent to the one given by [10, (3.43)].

(2) As for the defining relations of $U(\chi)$ for $\chi$ of Theorem 18 below, see [1], [2], [16], [17], [18], [20].

Using $\tilde{\Omega}$, we have the $\mathbb{C}$-algebra automorphism $\Omega : U \to U$ defined by $\Omega(K_\alpha) := K_{-\alpha}$, $\Omega(L_\alpha) := L_{-\alpha}$, $\Omega(E_i) := F_i L_{-\alpha_i}$, and $\Omega(F_i) := K_{-\alpha_i} E_i$.

Let $\pi : \tilde{U} \to U$ be the canonical map. We denote $\pi(\tilde{K}_\alpha)$, $\pi(\tilde{L}_\alpha)$, $\pi(\tilde{E}_i)$, $\pi(\tilde{F}_i)$, $\pi(\tilde{E}^\pm_{m,\alpha_i,\alpha_j})$, $\pi(\tilde{F}^\pm_{m,\alpha_i,\alpha_j})$ by $K_\alpha$, $L_\alpha$, $E_i$, $F_i$, $E^\pm_{m,\alpha_i,\alpha_j}$, $F^\pm_{m,\alpha_i,\alpha_j}$ respectively. Let $U^0 = U^0(\chi) := \pi(\tilde{U}^0)$, $U^\pm = U^\pm(\chi) := \pi(\tilde{U}^\pm)$. For $\alpha \in \mathbb{Z}\Pi$, let $U_\alpha = U(\chi)_\alpha := \pi(\tilde{U}_\alpha)$, and $U^\pm_\alpha = U^\pm(\chi)_\alpha := \pi(\tilde{U}^\pm_\alpha)$.

We can easily see

**Lemma 3.** *There exists a unique $\mathbb{C}$-linear isomorphism from $U^- \otimes U^0 \otimes U^+$ to $U$ sending $Y \otimes Z \otimes X$ to $YZX$ ($X \in U^+$, $Z \in U^0$, $Y \in U^-$). Moreover, we have $U^0 = \oplus_{\alpha,\beta\in\mathbb{Z}\Pi}\mathbb{C}K_\alpha L_\beta$, $U = \oplus_{\alpha\in\mathbb{Z}\Pi}U_\alpha$, $U^\pm = \oplus_{\alpha\in\pm\mathbb{Z}_{\geq 0}\Pi}U^\pm_\alpha$, and $\dim U^+_\alpha = \dim U^-_{-\alpha}$ for all $\alpha \in \mathbb{Z}_{\geq 0}\Pi$.*

## 1.4 Kharchenko-PBW theorem

Define the map $h^\chi : \mathbb{Z}\Pi \to \mathbb{N} \cup \{\infty\}$ by

(1.17)

$$h^\chi(\alpha) := \begin{cases} \infty & \text{if } (m)_{\chi(\alpha,\alpha)}! \neq 0 \text{ for all } m \in \mathbb{N}, \\ \mathrm{Max}\{m \in \mathbb{N} \,|\, (m)_{\chi(\alpha,\alpha)}! \neq 0\} & \text{otherwise.} \end{cases}$$

For $i$, $j \in I$, define $c^\chi_{ij} \in \{2\} \cup J_{-\infty,0} \cup \{-\infty\}$ by

$$c^\chi_{ij} := \begin{cases} 2 \text{ if } i = j, \\ -\infty \text{ if } i \neq j \text{ and } (m)_{q_{ii}}!(m; q_{ii}, q_{ij}q_{ji})! \neq 0 \text{ for all } m \in \mathbb{N}, \\ -\mathrm{Max}\{m \in \mathbb{Z}_{\geq 0} \,|\, (m)_{q_{ii}}!(m; q_{ii}, q_{ij}q_{ji})! = 0\} \text{ otherwise.} \end{cases} \tag{1.18}$$

By (1.9), we have

**Lemma 4.** *Let $i \in I$. Let $m \in \mathbb{Z}_{\geq 0}$. Then $F^m_i = 0$ if and only if $m > h^\chi(\alpha_i)$. In particular, if $m \in J_{0,h^\chi(\alpha_i)}$, then $\dim U^-_{-m\alpha_i} = 1$, and if $m > h^\chi(\alpha_i)$, then $\dim U^-_{-m\alpha_i} = 0$.*

By (1.9), (1.10) and (1.12), we have

**Lemma 5.** *Let $i$, $j \in I$ with $i \neq j$. Let $m \in \mathbb{Z}_{\geq 0}$.*

*(1) $F^+_{m,\alpha_i,\alpha_j} \neq 0$ if and only if $m \in J_{0,-c^\chi_{ij}}$.*

*(2) Let*

$$\mathcal{Y}_{i,j,m} := \{(m_1, m_2) \in \mathbb{Z}^2 \,|\, m_1 \in J_{0,-c^\chi_{ij}}, m_2 \in J_{0,h^\chi(\alpha_i)}, m_1 + m_2 = m\}. \tag{1.19}$$

*Then the elements $F^+_{m_1,\alpha_i,\alpha_j}F^{m_2}_i$ with $(m_1, m_2) \in \mathcal{Y}_{i,j,m}$ form a $\mathbb{C}$-basis of $U^-_{-m\alpha_i-\alpha_j}$.*

*(3) $\dim U^-_{-\alpha_i-\alpha_j} \in J_{1,2}$. Moreover, $q_{ij}q_{ji} = 1 \Leftrightarrow c^\chi_{ij} = 0 \Leftrightarrow \dim U^-_{-\alpha_i-\alpha_j} = 1 \Leftrightarrow F_iF_j = q_{ij}F_jF_i$.*

By the celebrated Kharchenko's PBW theorem, we have

**Theorem 6.** *(Kharchenko[12], see also [6, Section 3, (P)], and [10, (2.15)])*
*(1) There exists a unique pair*

$$(R^+ = R^+(\chi), \varphi = \varphi_\chi), \tag{1.20}$$

*where $R^+$ is a subset of $\mathbb{Z}_{\geq 0}\Pi \setminus \{0\}$, and $\varphi$ is a map from $R^+$ to $\mathbb{N}$, satisfying the condition that there exist $k \in \mathbb{N} \cup \{\infty\}$, a surjective map $\psi : J_{1,k} \to R^+$ with $|\psi^{-1}(\alpha)| = \varphi(\alpha)$ $(\alpha \in R^+)$, and $F[r] \in U^-_{-\psi(r)} \setminus \{0\}$ $(r \in J_{1,k})$ such that the elements*

$$F[1]^{x_1} \cdots F[m]^{x_m} \quad (m \in J_{1,k},\, x_y \in J_{0,h^\chi(\psi(y))}\, (y \in J_{1,m})) \tag{1.21}$$

*form a $\mathbb{C}$-basis of $U^-$ (where we mean that for $m \leq \bar{m}$, $F[1]^{x_1} \cdots F[m]^{x_m} = F[1]^{\bar{x}_1} \cdots F[\bar{m}]^{\bar{x}_{\bar{m}}}$ if and only if $x_y = \bar{x}_y$ for all $y \in J_{1,m}$ and $\bar{x}_{\bar{y}} = 0$ for all $\bar{y} \in J_{m+1,\bar{m}}$).*
*(2) (cf. [10, Theorem 4.5]) If $|R^+| < \infty$, then $\varphi(R^+) = \{1\}$.*

Once we know Theorem 6, the following lemma is clear from Lemmas 4 and 5.

**Lemma 7.** (1) $\Pi \subset R^+$.
(2) *For $i$, $j \in I$ with $i \neq j$, we have*

$$R^+ \cap (\alpha_j + \mathbb{Z}_{\geq 0}\alpha_i) = \{\alpha_j + n\alpha_i \,|\, n \in J_{0,-c^\chi_{ij}}\}, \tag{1.22}$$

*and $\varphi(\alpha_j + r\alpha_i) = 1$ for all $r \in J_{0,-c_{ij}}$.*
(3) *Let $I'$ be a non-empty proper subset of $I$. Let $I'' = I \setminus I'$. Then $R^+ = (R^+ \cap \oplus_{i \in I'}\mathbb{Z}_{\geq 0}\alpha_i) \uplus (R^+ \cap \oplus_{j \in I''}\mathbb{Z}_{\geq 0}\alpha_j)$ if and only if $q_{ij}q_{ji} = 1$ for all $i \in I'$ and all $j \in I''$.*

For $\chi \in \mathcal{X}_\Pi$, we say that $\chi$ is *irreducible* if its Dynkin diagram is connected, that is, for any two $i$, $j \in I$ with $i \neq j$, there exists $m \in \mathbb{Z}_{\geq 0}$, $i_r \in I$ $(r \in J_{1,m})$ such that $q_{i_t,i_{t+1}}q_{i_{t+1},i_t} \neq 1$ for all $t \in J_{0,m}$, where we let $i_0 := i$ and $i_{m+1} := j$.
Let $\mathcal{X}^{\mathrm{irr}}_\Pi := \{\chi \in \mathcal{X}_\Pi \,|\, \chi \text{ is irreducible}\}$.

## 1.5 Irreducible highest weight module

Let $\Lambda \in \mathrm{Ch}(U^0)$. By Lemma 3, we have a unique left $U$-module $M(\Lambda) = M_\chi(\Lambda)$ satisfying the following conditions.

(i) There exists $\tilde{v}_\lambda \in M_\chi(\Lambda) \setminus \{0\}$ such that $Z\tilde{v}_\Lambda = \Lambda(Z)\tilde{v}_\Lambda$ for all $Z \in U^0$ and $E_i\tilde{v}_\Lambda = 0$ for all $i \in I$.
(ii) The $\mathbb{C}$-linear map $U^- \to M(\Lambda)$, $Y \to Y\tilde{v}_\Lambda$, is bijective.

For $i \in I$ and $m \in \mathbb{Z}_{\geq 0}$, by (1.9), we have

$$E_i F_i^m \tilde{v}_\Lambda = (m)_{q_{ii}}(-q_{ii}^{1-m}\Lambda(K_{\alpha_i}) + \Lambda(L_{\alpha_i}))F_i^{m-1}\tilde{v}_\Lambda. \tag{1.23}$$

Let $\mathcal{N}$ be the maximal left submodule of $M(\Lambda)$ with $\mathcal{N} \subset \oplus_{\alpha \in \mathbb{Z}_{\geq 0}\Pi \setminus \{0\}} U^-_{-\alpha}\tilde{v}_\Lambda$. Then $\mathcal{N} = \oplus_{\alpha \in \mathbb{Z}_{\geq 0}\Pi \setminus \{0\}}(U^-_{-\alpha}\tilde{v}_\Lambda \cap \mathcal{N})$. Let $L(\Lambda) = L_\chi(\Lambda)$ be the quotient left $U$-module defined by

$$L(\Lambda) = L_\chi(\Lambda) := M(\Lambda)/\mathcal{N}. \tag{1.24}$$

We denote the element $\tilde{v}_\Lambda + \mathcal{N}$ of $L(\Lambda)$ by $v_\Lambda$. For $\alpha \in \mathbb{Z}\Pi$, let $L(\Lambda)_\alpha = L_\chi(\Lambda)_\alpha := U^-_\alpha v_\Lambda$. Then $L(\Lambda) = \oplus_{\alpha \in \mathbb{Z}_{\geq 0}\Pi} L(\Lambda)_{-\alpha}$. We also have $L(\Lambda)_0 = \mathbb{C}v_\Lambda$, and $\dim L(\Lambda)_0 = 1$.

By (1.23), for $m \in \mathbb{N}$, we have

$$F_i^m v_\Lambda = 0 \quad \Leftrightarrow \quad (m)_{q_{ii}}!(m; q_{ii}^{-1}, \Lambda(K_{\alpha_i}L_{-\alpha_i}))! = 0. \tag{1.25}$$

Let

$$h_{\chi,\Lambda,i} := \begin{cases} \infty \quad \text{if } (m)_{q_{ii}}!(m; q_{ii}^{-1}, \Lambda(K_{\alpha_i}L_{-\alpha_i}))! \neq 0 \text{ for all } m \in \mathbb{N}, \\ \text{Max}\{m \in \mathbb{Z}_{\geq 0} | (m)_{q_{ii}}!(m; q_{ii}^{-1}, \Lambda(K_{\alpha_i}L_{-\alpha_i}))! \neq 0\} \\ \quad \text{otherwise.} \end{cases} \tag{1.26}$$

By (1.25), we have

$$\dim L(\Lambda) < \infty \Rightarrow \forall i \in I,\ h_{\chi,\Lambda,i} < \infty. \tag{1.27}$$

## 1.6 Notation $\equiv$

For $\chi \in \mathcal{X}_\Pi$, define the map $\kappa_\chi : \mathrm{Ch}(U^0(\chi)) \to (\mathbb{C}^\times)^N$ by $\kappa_\chi(\Lambda) := (\Lambda(K_{\alpha_i}L_{-\alpha_i}))_{i \in I}$.

*Notation* 2. Let $\chi$, $\chi' \in \mathcal{X}_\Pi$. Let $q_{ij} := \chi(\alpha_i, \alpha_j)$ and $q'_{ij} := \chi'(\alpha_i, \alpha_j)$. We write

$$\chi \equiv \chi' \tag{1.28}$$

if $q_{ii} = q'_{ii}$ for all $i \in I$ and $q_{jk}q_{kj} = q'_{jk}q'_{kj}$ for all $j$, $k \in I$. By (1.18),

$$\chi \equiv \chi' \quad \Rightarrow \quad \forall i, \forall j \in I,\ c^\chi_{ij} = c^{\chi'}_{ij}. \tag{1.29}$$

Let $\Lambda \in \mathrm{Ch}(U^0(\chi))$ and $\Lambda' \in \mathrm{Ch}(U^0(\chi'))$. We write

$$(\chi, \Lambda) \equiv (\chi', \Lambda') \tag{1.30}$$

if $\chi \equiv \chi'$ and $\kappa_\chi(\Lambda) = \kappa_{\chi'}(\Lambda')$.

Note that

$$(\chi, \Lambda) \equiv (\chi', \Lambda') \quad \Longrightarrow \quad \forall i \in I,\ h_{\chi,\Lambda,i} = h_{\chi',\Lambda',i}. \tag{1.31}$$

# 2 Some finite-type bi-homomorphisms

## 2.1 Finite-type bi-homomorphisms

Let $\mathcal{X}_\Pi^{\mathrm{fin}} := \{\chi \in \mathcal{X}_\Pi \,|\, |R^+(\chi)| < \infty\}$. If $\chi \in \mathcal{X}_\Pi^{\mathrm{fin}} \cap \mathcal{X}_\Pi^{\mathrm{irr}}$, we say that $\chi$ is *finite-type*.

Let $\mathcal{X}_\Pi^{\prime\prime,\mathrm{fin}} := \{\chi \in \mathcal{X}_\Pi \,|\, \forall i, \forall j,\, c_{ij}^\chi \neq -\infty\,\}$. By (1.22), we have

$$\mathcal{X}_\Pi^{\mathrm{fin}} \subset \mathcal{X}_\Pi^{\prime\prime,\mathrm{fin}}. \tag{2.1}$$

Let $\chi \in \mathcal{X}_\Pi^{\prime\prime,\mathrm{fin}}$. Let $i \in I$. Define $s_i = s_i^\chi \in \mathrm{Aut}_{\mathbb{Z}}(\mathbb{Z}\Pi)$ by $s_i(\alpha_j) = \alpha_j - c_{ij}\alpha_i$. Note that $s_i(\alpha_i) = -\alpha_i$, and $s_i^2 = \mathrm{id}_{\mathbb{Z}\Pi}$ $(i \in I)$. Define $i \triangleright \chi \in \mathcal{X}_\Pi$ by

$$i \triangleright \chi(\alpha, \beta) := \chi(s_i(\alpha), s_i(\beta)) \tag{2.2}$$

for all $\alpha$, $\beta \in \mathbb{Z}\Pi$.

*Notation* 3. Let $\chi$, $\chi' \in \mathcal{X}_\Pi^{\prime\prime,\mathrm{fin}}$. We write $\chi \sim \chi'$ if there exist $m \in \mathbb{N}$, $i_t \in I$ $(t \in J_{1,m})$ and $\chi_r \in \mathcal{X}_\Pi^{\prime\prime,\mathrm{fin}}$ $(r \in J_{1,m+1})$ such that $\chi = \chi_1$, $\chi' = \chi_{m+1}$, and $i_t \triangleright \chi_t = \chi_{t+1}$ $(t \in J_{1,m})$.

Let $\mathcal{X}_\Pi^{\prime,\mathrm{fin}} := \{\chi \in \mathcal{X}_\Pi^{\prime\prime,\mathrm{fin}} \,|\, \forall \chi' \sim \chi,\, \forall i \in I,\, i \triangleright \chi' \in \mathcal{X}_\Pi^{\prime\prime,\mathrm{fin}}\,\}$.

Let $R(\chi) := R^+(\chi) \cup (-R^+(\chi))$ for $\chi \in \mathcal{X}_\Pi$. It is well-known that

**Theorem 8.** ([6, Proposition 1], *see also* [9, Example 4]) *Let* $\chi \in \mathcal{X}_\Pi^{\prime\prime,\mathrm{fin}}$.

(1) *We have*

$$s_i^\chi(R^+(\chi) \setminus \{\alpha_i\}) = R^+(i \triangleright \chi) \setminus \{\alpha_i\}. \tag{2.3}$$

*In particular,*

$$s_i^\chi(R(\chi)) = R(i \triangleright \chi). \tag{2.4}$$

*and*

$$\chi \in \mathcal{X}_\Pi^{\mathrm{fin}} \implies i \triangleright \chi \in \mathcal{X}_\Pi^{\mathrm{fin}}. \tag{2.5}$$

(2) *For* $\chi \in \mathcal{X}_\Pi^{\prime,\mathrm{fin}}$, $\{(R(\chi'), \Pi, (s_i^{\chi'})_{i\in I}) | \chi' \sim \chi\}$ *satisfies the axioms of [9, Definition 2].*

By (2.1) and (2.5), we have

$$\mathcal{X}_\Pi^{\mathrm{fin}} \subset \mathcal{X}_\Pi^{\prime,\mathrm{fin}}. \tag{2.6}$$

Let $\chi \in \mathcal{X}_\Pi^{\prime,\mathrm{fin}}$. By (2.3) and (1.22), we have $c_{ij}^{i\triangleright\chi} = c_{ij}^\chi$. Hence $s_i^{i\triangleright\chi} = s_i^\chi$, and we have

$$i \triangleright i \triangleright \chi = \chi. \tag{2.7}$$

The following lemma is an easy exercise for the reader.

**Lemma 9.** *Let* $\chi \in \mathcal{X}_\Pi^{\prime\prime,\mathrm{fin}}$. *Let* $i \in I$ *be such that* $\chi(\alpha_i, \alpha_i) \in P_\infty$. *Then* $i \triangleright \chi \equiv \chi$.

Let $\chi$, $\chi' \in \mathcal{X}_\Pi^{\prime\prime,\mathrm{fin}}$. Let $i \in I$. By (1.29), we have

$$\chi \equiv \chi' \implies s_i^\chi = s_i^{\chi'},\; i \triangleright \chi \equiv i \triangleright \chi'. \tag{2.8}$$

## 2.2 Longest element of Weyl groupoid associated with a finite-type bi-homomorphism

For $n \in \mathbb{N} \cup \{\infty\}$, let $\Xi_n$ be the set of maps from $J_{1,n}$ to $I$.

*Notation* 4. Let $\chi \in \mathcal{X}'^{,\mathrm{fin}}_{\Pi}$. Let $n \in \mathbb{N} \cup \{\infty\}$, and let $f \in \Xi_n$. For $t \in J_{0,n}$, define $\chi_{f,t} \in \mathcal{X}'^{,\mathrm{fin}}_{\Pi}$ and $1^{\chi}s_{f,t} \in \mathrm{Aut}_{\mathbb{Z}}(\mathbb{Z}\Pi)$ inductively as follows. Let $\chi_{f,0} := \chi$, and let $1^{\chi}s_{f,0} := \mathrm{id}_{\mathbb{Z}\Pi}$. If $t \in J_{1,n}$, let $\chi_{f,t} := f(t) \triangleright \chi_{f,t-1}$, and let $1^{\chi}s_{f,t} := 1^{\chi}s_{f,t-1}s^{\chi_{f,t}}_{f(t)}$.

Let $\chi \in \mathcal{X}'^{,\mathrm{fin}}_{\Pi}$, $f \in \Xi_\infty$ and $t \in \mathbb{Z}_{\geq 0}$. Let $w := 1^{\chi}s_{f,t}$. Let

$$\ell(w) := \min\{l \in \mathbb{Z}_{\geq 0} \,|\, \exists f' \in \Xi_\infty, \text{ s.t. } \chi_{f',l} = \chi_{f,t},\ 1^{\chi}s_{f',l} = w\}. \tag{2.9}$$

By Theorem 8 and [9, Lemma 8 (iii)], we have

$$\ell(w) = |w(R^+(\chi_{f,t})) \cap -\mathbb{Z}_{\geq 0}\Pi|. \tag{2.10}$$

Hence we have

$$\ell(1^{\chi}s_{f,t+1}) = \begin{cases} \ell(w)+1 & \text{if } w(\alpha_{f(t+1)}) \in R^+(\chi), \\ \ell(w)-1 & \text{if } w(\alpha_{f(t+1)}) \in -R^+(\chi). \end{cases} \tag{2.11}$$

For $f_1$, $f_2 \in \Xi_\infty$ and $l_1$, $l_2 \in \mathbb{Z}_{\geq 0}$ with $1^{\chi}s_{f_1,l_1} = 1^{\chi}s_{f_2,l_2}$, we have $\chi_{f_1,l_1} = \chi_{f_2,l_2}$ since $\ell((1^{\chi}s_{f_1,l_1})^{-1}1^{\chi}s_{f_2,l_2}) = 0$. Hence $\ell(w) = \min\{l \in \mathbb{Z}_{\geq 0} \,|\, \exists f' \in \Xi_\infty,\ 1^{\chi}s_{f',l} = w\}$.

**Lemma 10.** *Let $\chi \in \mathcal{X}^{\mathrm{fin}}_{\Pi}$, $n := |R^+(\chi)|$, $f \in \Xi_n$, and $t \in J_{1,n}$. Assume that $1^{\chi}s_{f,y}(\alpha_{f(y+1)}) \in R^+(\chi)$ hold for all $y \in J_{1,t-1}$. Then we have:*

(1) *$\ell(1^{\chi}s_{f,t}) = t$.*

(2) *Assume $t \in J_{1,n-1}$, Then there exists $i \in I$ such that $1^{\chi}s_{f,t}(\alpha_i) \in R^+(\chi)$. Moreover, if $1^{\chi}s_{f,t}(\alpha_{f(t+1)}) \in -R^+(\chi)$, then $\ell(1^{\chi}s_{f,t+1}) = t-1$.*

(3) *Assume $t = n$. Then we have*

$$1^{\chi}s_{f,n}(\Pi) = -\Pi, \tag{2.12}$$

*and*

$$R^+(\chi) = \{\, 1^{\chi}s_{f,r-1}(\alpha_{f(r)}) \,|\, r \in J_{1,n} \,\}, \tag{2.13}$$

*and for $n' \in \mathbb{N}$ and $f' \in \Xi_{n'}$.*

(4) *For $\chi' \in \mathcal{X}_{\Pi}$ with $\chi' \equiv \chi$, we have $\chi' \in \mathcal{X}^{\mathrm{fin}}_{\Pi}$ and $R(\chi') = R(\chi)$.*

*Proof.* (1), (2) and (2.12) are clear from (2.11).

We show (2.13). By (1), we see that for all $t_1$, $t_2 \in J_{0,n}$,

$$t_1 \leq t_2 \ \Rightarrow\ \ell((1^{\chi}s_{f,t_1})^{-1}1^{\chi}s_{f,t_2}) = t_2 - t_1. \tag{2.14}$$

Assume that $1^\chi s_{f,r_1-1}(\alpha_{f(r_1)}) = 1^\chi s_{f,r_2-1}(\alpha_{f(r_2)})$ for some $r_1$, $r_2 \in J_{1,n}$ with $r_1 < r_2$. Applying $(1^\chi s_{f,r_1})^{-1}$, we have $-\alpha_{f(r_1)} = (1^\chi s_{f,r_1})^{-1} 1^\chi s_{f,r_2-1}(\alpha_{f(r_2)})$. By (2.11), $\ell((1^\chi s_{f,r_1})^{-1} 1^\chi s_{f,r_2}) = \ell((1^\chi s_{f,r_1})^{-1} 1^\chi s_{f,r_2-1}) - 1$, which contradicts (2.14). Hence we have (2.13), as desired.

We show (4). By (2.8), we have $\chi' \in \mathcal{X}_\Pi^{\prime,\mathrm{fin}}$ and $1^{\chi'} s_{f,t} = 1^\chi s_{f,t}$ $(t \in J_{0,n})$. By (2.10) and (2.12), we have $\chi' \in \mathcal{X}_\Pi^{\mathrm{fin}}$ and $|R^+(\chi')| \leq n$. By (2.13), we have $R^+(\chi) = R^+(\chi')$. This completes the proof. □

**Lemma 11.** *Let $\chi \in \mathcal{X}_\Pi^{\mathrm{fin}}$. Then $\dim U^-(\chi) < \infty$ if and only if $\chi(\alpha_i, \alpha_i) \notin P_\infty$ for all $i$, $j \in I$.*

*Proof.* By (2.13), (2.4), and Theorem 6, we see that $\dim U^-(\chi) < \infty$ if and only if $\chi'(\alpha_i, \alpha_i) \notin P_\infty$ for all $i \in I$, and all $\chi' \in \mathcal{X}_\Pi^{\mathrm{fin}}$ with $\chi' \sim \chi$. Then we can easily see that this lemma holds. □

## 2.3 Classification of irreducible infinite-dimensional finite-type bi-homomorphisms

Let $n \in \mathbb{N}$. Let $\mathbb{Z}^n$ be a rank-$n$ free $\mathbb{Z}$-module with a basis $\{e_r | r \in J_{1,n}\}$. Let $m \in J_{0,n}$. Define the map $\eta_{m|n-m} : \mathbb{Z}^n \times \mathbb{Z}^n \to \mathbb{Z}$ by $\eta_{m|n-m}(\sum_{r=1}^n x_r e_r, \sum_{r'=1}^n x'_{r'} e_{r'}) := \sum_{l=1}^m x_l x'_l - \sum_{l'=m+1}^n x_{l'} x'_{l'}$. Define the map $p_{m|n-m} : \mathbb{Z}^n \to J_{0,1}$ by $p_{m|n-m}(z) := \delta_{0,\eta_{m|n-m}(z,z)}$.

For $\chi \in \mathcal{X}_\Pi$, and $\sigma \in \mathfrak{S}(I)$, define $\chi^\sigma \in \mathcal{X}_\Pi$ by $\chi^\sigma(\alpha_i, \alpha_j) := \chi(\alpha_{\sigma^{-1}(i)}, \alpha_{\sigma^{-1}(j)})$ for all $i$, $j \in I$.

By the Heckenberger's classification [7, Tables 1-4, Theorems 17, 22] and Lemma 11, we have

**Theorem 12.** ([7]) *For $b \in J_{1,10}$, let $\mathcal{X}_\Pi^{(b)}$ be the set of $\chi \in \mathcal{X}_\Pi$ given in $(b)$ below, and let $\mathcal{X}_\Pi^{(b),\prime} := \{\, \chi' \mid \exists \chi \in \mathcal{X}_\Pi^{(b)}, \exists \sigma \in \mathfrak{S}(I), \chi' \sim \chi^\sigma \,\}$. Then we have*

$$\{\chi \in \mathcal{X}_\Pi^{\mathrm{irr}} \cap \mathcal{X}_\Pi^{\mathrm{fin}} \mid \dim U^-(\chi) = \infty \,\} = \biguplus_{b \in J_{1,10}} \mathcal{X}_\Pi^{(b),\prime}. \tag{2.15}$$

*In the following, let $q_{ij} := \chi(\alpha_i, \alpha_j)$.*

(1) ($A_N$ $(N \in \mathbb{N})$, $B_N$ $(N \in J_{2,\infty})$, $C_N$ $(N \in J_{3,\infty})$, $D_N$ $(N \in J_{4,\infty})$, $E_N$ $(N \in J_{6,8})$, $F_4$ $(N = 4)$, $G_2$ $(N = 2)$, $B(0,N)$ $(N \in \mathbb{N})$) *There exist $N \times N$ irreducible finite-type Cartan matrix $A = [a_{ij}]_{i,j \in I}$, $d_i \in \mathbb{N}$ $(i \in I)$, and $\hat{q} \in P_\infty$ such that $d_i a_{ij} = d_j a_{ji}$, $q_{ii} = \hat{q}^{2d_i}$ and $q_{ij} q_{ji} = \hat{q}^{2d_i a_{ij}}$ hold for all $i$, $j \in I$.*

(2) ($sl(m+1|N-m)$, *or* [7, Table 4, Row 2]) *Assume $N \in J_{2,\infty}$ and let $m \in J_{0,\frac{N-1}{2}}$. Let $\bar{\alpha}_i := e_i - e_{i+1} \in \mathbb{Z}^{N+1}$ for $i \in I$. There exists $\hat{q} \in P_\infty$ such that $q_{ii} = (-1)^{p_{m+1|N-m}(\bar{\alpha}_i)} \hat{q}^{\eta_{m+1|N-m}(\bar{\alpha}_i, \bar{\alpha}_i)}$ and $q_{ij} q_{ji} = \hat{q}^{2\eta_{m+1|N-m}(\bar{\alpha}_i, \bar{\alpha}_j)}$ hold for all $i$, $j \in I$.*

(3) ($B(m, N-m)$, *or* [7, Table 4, Row 4]) *Assume $N \in J_{2,\infty}$ and let $m \in J_{1,N-1}$. Let $\bar{\alpha}_i := e_i - e_{i+1} \in \mathbb{Z}^N$ for $i \in J_{1,N-1}$. Let $\bar{\alpha}_N := e_N \in \mathbb{Z}^N$. There exists*

$\hat{q} \in P_\infty$ *such that* $q_{ii} = (-1)^{p_{N-m|m}(\bar{\alpha}_i)}\hat{q}^{\eta_{N-m|m}(\bar{\alpha}_i,\bar{\alpha}_i)}$ *and* $q_{ij}q_{ji} = \hat{q}^{2\eta_{N-m|m}(\bar{\alpha}_i,\bar{\alpha}_j)}$ *hold for all* $i$, $j \in I$.

(4) ($C(N)$, *or* [7, Table 4, Row 9]) *Assume* $N \in J_{3,\infty}$. *Let* $\bar{\alpha}_i := e_i - e_{i+1} \in \mathbb{Z}^N$ *for* $i \in J_{1,N-1}$. *Let* $\bar{\alpha}_N := 2e_N \in \mathbb{Z}^N$. *There exists* $\hat{q} \in P_\infty$ *such that* $q_{ii} = (-1)^{p_{1|N-1}(\bar{\alpha}_i)}\hat{q}^{\eta_{1|N-1}(\bar{\alpha}_i,\bar{\alpha}_i)}$ *and* $q_{ij}q_{ji} = \hat{q}^{2\eta_{1|N-1}(\bar{\alpha}_i,\bar{\alpha}_j)}$ *hold for all* $i$, $j \in I$.

(5) ($D(m, N-m)$, *or* [7, Table 4, Row 10]) *Assume* $N \in J_{4,\infty}$ *and let* $m \in J_{2,N-1}$. *Let* $\bar{\alpha}_i := e_i - e_{i+1} \in \mathbb{Z}^N$ *for* $i \in J_{1,N-1}$. *Let* $\bar{\alpha}_N := e_{N-1} + e_N \in \mathbb{Z}^N$. *There exists* $\hat{q} \in P_\infty$ *such that* $q_{ii} = (-1)^{p_{N-m|m}(\bar{\alpha}_i)}\hat{q}^{\eta_{N-m|m}(\bar{\alpha}_i,\bar{\alpha}_i)}$ *and* $q_{ij}q_{ji} = \hat{q}^{2\eta_{N-m|m}(\bar{\alpha}_i,\bar{\alpha}_j)}$ *hold for all* $i$, $j \in I$.

(6) ($F(4)$, *or* [7, Table 3, Row 9]) *Assume* $N = 4$. *The Dynkin diagram of* $\chi$ *is the one of* $\chi_{f,0}$ *in Figure 6 with* $\hat{q} \in P_\infty$.

(7) ($G(3)$, *or* [7, Table 2, Row 7]) *Assume* $N = 3$. *The Dynkin diagram of* $\chi$ *is the one of* $\chi_{f,0}$ *in Figure 7 with* $\hat{q} \in P_\infty$.

(8) ($D(2,1;x)$, *or* [7, Table 2, Row 9]) *Assume* $N = 3$. *The Dynkin diagram of* $\chi$ *is the one of* $\chi_{f,0}$ *in Figure 8 with* $\hat{q} \in P_\infty$ *and* $\hat{r} \in \mathbb{C}^\times \setminus \{1, \hat{q}^{-1}\}$.

(9) ([7, Table 1, Row 5]) *Assume* $N = 2$. *The Dynkin diagram of* $\chi$ *is the one of* $\chi_{f,0}$ *in Figure 9 with* $\hat{q} \in P_\infty$ *and* $\hat{\zeta} \in P_3$.

(10) ([7, Table 3, Row 14]) *Assume* $N = 4$. *The Dynkin diagram of* $\chi$ *is the one of* $\chi_{f,0}$ *in Figure 10 with* $\hat{q} \in P_\infty$.

## 2.4 Longest elements of the classical Weyl groups

In the proof of Theorem 18 below, we use the notation and facts as follows. Assume $b \in J_{2,5}$, and keep the notation in Theorem 12 (b). Let $N' := N + \delta_{b2}$. For $z \in \mathbb{Z}^{N'} \setminus \{0\}$, define $\bar{s}_z \in \mathrm{Aut}_{\mathbb{Z}}(\mathbb{Z}^{N'})$ by $\bar{s}_i(y) := y - \frac{2\eta_{0|N'}(y,z)}{\eta_{0|N'}(z,z)}z$. Let $\bar{s}_i := \bar{s}_{\bar{\alpha}_i}$ ($i \in I$). Let $\bar{W}$ be the subgroup of $\mathrm{Aut}_{\mathbb{Z}}(\mathbb{Z}^{N'})$ generated by $\bar{s}_i$ ($i \in I$). Let $\bar{X} := \{\bar{s}_i | i \in I\}$. Then $(\bar{W}, \bar{X})$ is the Coxeter system of type $A_N$, $B_N$, $C_N (= B_N)$, $D_N$ for $b = 2$, 3, 4, 5 respectively. Let $\bar{w}_0$ be the longest element of $(\bar{W}, \bar{X})$. Then we have

$$\bar{w}_0(e_y) = \begin{cases} e_{N+2-y} & b = 2, \\ e_N & b = 5,\ y = N \in 2\mathbb{Z}, \\ -e_y & \text{otherwise} \end{cases} \tag{2.16}$$

($y \in J_{1,N'}$), and we have a reduced expression of $\bar{w}_0$ as follows:

$$\bar{w}_0 = \begin{cases} (\bar{s}_1\bar{s}_2\cdots\bar{s}_N)(\bar{s}_1\bar{s}_2\cdots\bar{s}_{N-1})\cdots(\bar{s}_1\bar{s}_2)\bar{s}_1 & b = 2, \\ (\bar{s}_1\bar{s}_2\cdots\bar{s}_N)^N & b \in J_{3,4}, \\ (\bar{s}_1\cdots\bar{s}_{N-2}\bar{s}_{N_1})(\bar{s}_1\cdots\bar{s}_{N-2}\bar{s}_{N_2})\cdots(\bar{s}_1\cdots\bar{s}_{N-2}\bar{s}_{N_N}) & b = 5. \end{cases} \tag{2.17}$$

where $N_z := N - \delta_{(-1)^z,-1}$.

For a non-empty subset $I'$ of $I$, let $\bar{X}^{I'} := \{\bar{s}_i | i \in I'\}$, and let $\bar{W}^{I'}$ be the subgroup of $\bar{W}$ generated by $\bar{X}^{I'}$. Then $(\bar{W}^{I'}, \bar{X}^{I'})$ is also a Coxeter system.

Assume $b = 2$. Let $I'' := I \setminus \{m+1\}$. Using the same results for $(\bar{W}^{I''}, \bar{X}^{I''})$ as in (2.16)-(2.17), we can see that we have a reduced expression of $\bar{w}_0$ as follows.

$$
\begin{aligned}
\bar{w}_0 \;=\;& (\underbrace{\bar{s}_1\bar{s}_2\cdots\bar{s}_m}_{m})(\underbrace{\bar{s}_1\bar{s}_2\cdots\bar{s}_{m-1}}_{m-1})\cdots(\underbrace{\bar{s}_1\bar{s}_2}_{2})\underbrace{\bar{s}_1}_{1} \\
&\cdot(\underbrace{\bar{s}_{m+2}\bar{s}_{m+3}\cdots\bar{s}_N}_{N-m-1})(\underbrace{\bar{s}_{m+2}\bar{s}_{m+3}\cdots\bar{s}_{N-1}}_{N-m-2})\cdots(\underbrace{\bar{s}_{m+2}\bar{s}_{m+3}}_{2})\underbrace{\bar{s}_{m+2}}_{1} \\
&\cdot(\underbrace{\bar{s}_{m+1}\bar{s}_{m+2}\cdots\bar{s}_N}_{N-m})(\underbrace{\bar{s}_m\bar{s}_{m+1}\cdots\bar{s}_{N-1}}_{N-m})\cdots(\underbrace{\bar{s}_1\bar{s}_2\cdots\bar{s}_{N-m}}_{N-m}).
\end{aligned}
\tag{2.18}
$$

Assume $b = 3$. Let $\bar{X}' := \{\bar{s}_i | i \in J_{1,N-m-1}\}, \cup\{\bar{s}_{e_{N-m}}\}$. Let $\bar{W}'$ be the subgroup of $\bar{W}$ generated by $\bar{X}'$. Note that $(\bar{W}', \bar{X}')$ is the Coxeter system of type $B_{N-m}$. Note that $\bar{s}_{e_{N-m}} = \bar{s}_{N-m}\bar{s}_{N-m+1}\cdots\bar{s}_{N-1}\bar{s}_N\bar{s}_{N-1}\cdots\bar{s}_{N-m+1}\bar{s}_{N-m}$. Let $I''' := J_{N-m+1,N}$. Using the Coxeter system $(\bar{W}' \times \bar{W}^{I'''}, \bar{X}' \times \bar{X}^{I'''})$ of type $B_{N-m} \times B_m$, by (2.16)-(2.17), we can see that we have a reduced expression of $\bar{w}_0$ as follows.

$$
\begin{aligned}
\bar{w}_0 \;=\;& (\underbrace{\bar{s}_{N-m+1}\bar{s}_{N-m+2}\cdots\bar{s}_N}_{m})^m \\
&\cdot(\underbrace{\bar{s}_1\bar{s}_2\cdots\bar{s}_{N-1}\bar{s}_N\bar{s}_{N-1}\cdots\bar{s}_{N-m+1}\bar{s}_{N-m}}_{N+m})^{N-m}.
\end{aligned}
\tag{2.19}
$$

Assume $b = 4$. Similarly we have

$$
\bar{w}_0 = (\underbrace{\bar{s}_2\bar{s}_3\cdots\bar{s}_N}_{N-1})^{N-1} \cdot \underbrace{\bar{s}_1\bar{s}_2\cdots\bar{s}_{N-1}\bar{s}_N\bar{s}_{N-1}\cdots\bar{s}_2\bar{s}_1}_{2N-1}.
\tag{2.20}
$$

Assume $b = 5$. Let $\bar{s}'_N := \bar{s}_{e_N}$. Similarly we have

$$
\begin{aligned}
\bar{w}_0 \;=\;& \underbrace{(\bar{s}_{N-m+1}\cdots\bar{s}_{N-2}\bar{s}_{N_1})(\bar{s}_{N-m+1}\cdots\bar{s}_{N-2}\bar{s}_{N_2})\cdots(\bar{s}_{N-m+1}\cdots\bar{s}_{N-2}\bar{s}_{N_m})}_{m(m-1)} \\
&\cdot(\underbrace{\bar{s}_1\bar{s}_2\cdots\bar{s}_{N-1}\bar{s}'_N\bar{s}_{N-1}\cdots\bar{s}_{N-m+1}\bar{s}_{N-m}}_{N+m})^{N-m}.
\end{aligned}
\tag{2.21}
$$

*Notation* 5. For $b = 2$ (resp. 3, resp. 4, resp. 5), let $r_{(b)} := \frac{N(N+1)}{2}$ (resp. $N^2$, resp. $N^2$, resp. $N^2 - m$), and define $f_{(b)} \in \Xi_{r_{(b)}}$ in the way that for $x \in J_{1,r_{(b)}}$, the $x$-th factor of RHS of (2.18) (resp. (2.19), resp. (2.20), resp. (2.21)) is $\bar{s}_{f_{(b)}(x)}$ or $\bar{s}'_{f_{(b)}(x)}$.

# 3 Lusztig isomorphisms

In this section, fix $\chi \in \mathcal{X}_\Pi^{\mathrm{fin}}$, let $q_{ij} := \chi(\alpha_i, \alpha_j)$ $(i, j \in I)$, and fix $\Lambda \in \mathrm{Ch}(U^0(\chi))$.

**Theorem 13.** ([8]) *There exists a unique* $\mathbb{C}$*-algebra isomorphism*

$$T_i = T_i^{i \triangleright \chi} : U(i \triangleright \chi) \to U(\chi) \tag{3.1}$$

*such that*

$$\begin{array}{l} T_i(K_\alpha) = K_{s_i^{i\triangleright\chi}(\alpha)}, \quad T_i(L_\alpha) = L_{s_i^{i\triangleright\chi}(\alpha)}, \\ T_i(E_i) = F_i L_{-\alpha_i}, \quad T_i(F_i) = K_{-\alpha_i} E_i, \\ T_i(E_j) = E^+_{-c^\chi_{ij},\alpha_i,\alpha_j}, \\ T_i(F_j) = \frac{1}{(-c^\chi_{ij})_{q_{ii}}!(-c^\chi_{ij};q_{ii},q_{ij}q_{ji})!} F^+_{-c^\chi_{ij},\alpha_i,\alpha_j}. \end{array} \tag{3.2}$$

*hold for all* $\alpha \in \mathbb{Z}\Pi$ *and all* $j \in I \setminus \{i\}$.

**Lemma 14.** *Let* $i \in I$ *and* $h := h_{\chi,\Lambda,i}$. *Assume* $h \neq \infty$. *Then there exist a unique* $i \triangleright \Lambda \in \mathrm{Ch}(U^0(i \triangleright \chi))$ *and a unique* $\mathbb{C}$*-linear isomorphism*

$$\hat{T}_i = \hat{T}_i^{i\triangleright\chi, i\triangleright\Lambda} : L_{i\triangleright\chi}(i \triangleright \Lambda) \to L_\chi(\Lambda) \tag{3.3}$$

*such that*

$$\forall X \in U(i \triangleright \chi), \quad \hat{T}_i(X v_{i\triangleright\chi}) = T_i(X) F_i^h v_\Lambda. \tag{3.4}$$

*Moreover, we have*

$$h_{i\triangleright\chi, i\triangleright\Lambda, i} = h, \tag{3.5}$$

*and*

$$i \triangleright i \triangleright \Lambda = \Lambda. \tag{3.6}$$

*Further, for* $\chi' \in \mathcal{X}_\Pi^{\mathrm{fin}}$ *and* $\Lambda' \in \mathrm{Ch}(U^0(\chi'))$, *we have*

$$(\chi, \Lambda) \equiv (\chi', \Lambda') \implies (i \triangleright \chi, i \triangleright \Lambda) \equiv (i \triangleright \chi', i \triangleright \Lambda'). \tag{3.7}$$

*Proof.* We can regard $L_\chi(\Lambda)$ as a left $U(i \triangleright \chi)$-module defined by $X \cdot u := T_i(X)u$ ($X \in U(i \triangleright \chi)$, $u \in L_\chi(\Lambda)$). Then there exist $i \triangleright \chi \in \mathrm{Ch}(U^0(i \triangleright \chi))$, and a $U(i \triangleright \chi)$-module homomorphism $z : M_{i\triangleright\chi}(i \triangleright \Lambda) \to L_\chi(\Lambda)$ such that $z(X\tilde{v}_{i\triangleright\chi}) \to T_i(X)F_i^h v_\Lambda$ ($X \in U(i \triangleright \chi)$). We have

$$\forall k \in J_{0,h}, \ z^{-1}(\mathbb{C}F_i^k v_\Lambda) = \mathbb{C}F_i^{h-k}\tilde{v}_{i\triangleright\chi} \tag{3.8}$$

(cf. (1.23)). Since $z(E_i F_i^{h+1}\tilde{v}_{i\triangleright\chi}) = 0$, by (3.8), we have $E_i F_i^{h+1}\tilde{v}_{i\triangleright\chi} = 0$. Hence (3.5) holds. Hence we see that the linear isomorphism $\hat{T}_i$ of (3.3) exists. The rest of the statement is clear. This completes the proof. □

The following lemme is easy exercise for the reader, use also Lemma 9 and (1.31).


$2 \triangleright \Lambda$
$\Lambda$
$1 \triangleright 2 \triangleright \Lambda$
$1 \triangleright \Lambda$
$1 \triangleright 2 \triangleright 1 \triangleright \Lambda$
$= 2 \triangleright 1 \triangleright 2 \triangleright \Lambda$
$2 \triangleright 1 \triangleright \Lambda$

Figure 1: $L_\chi(\Lambda)$ for $\chi \in \mathcal{X}_\Pi^{(2)}$ with $N = 2$, $m = 1$, $\lambda_1 \neq -1$, $\lambda_2 = \hat{q}^8$, $\lambda_1\lambda_2\hat{q}^2 \neq -1$, where $\lambda_i := \Lambda(K_{\alpha_i} L_{-\alpha_i})$

**Lemma 15.** *Let* $i \in I$ *be such that* $q_{ii} \in P_\infty$ *and* $h_{\chi,\Lambda,i} \neq \infty$. *Then* $(i \triangleright \chi, i \triangleright \Lambda) \equiv (\chi, \Lambda)$, *and* $h_{i\triangleright\chi, i\triangleright\Lambda, j} = h_{\chi,\Lambda,j}$ *for all* $j \in I$.

*Notation* 6. Let $n \in \mathbb{N}$, and let $f \in \Xi_n$. Recall Notation 4. Let $\Lambda_{f,0} =: \Lambda$. Let $\Lambda_{f,t} =: f(t) \triangleright \Lambda_{f,t-1}$ for $t \in J_{1,n}$ such that $\forall r \in J_{0,t-1}$, $h_{\chi_{f,r},\Lambda_{f,r},f(r+1)} < \infty$. Let

$$H(\chi, \Lambda, f) := \mathrm{Max}\{t \in J_{0,n} \,|\, \forall r \in J_{0,t-1},\ h_{\chi_{f,r},\Lambda_{f,r},f(r+1)} < \infty\}. \tag{3.9}$$

Then the following lemma is clear from (1.27) and Lemma 14.

**Lemma 16.** *Let* $n \in \mathbb{N}$, *and let* $f \in \Xi_n$. *Assume* $1^\chi s_{f,n}(\Pi) = -\Pi$. *Then* $\dim L_\chi(\Lambda) < \infty$ *if and only if* $H(\chi, \Lambda, f) = n$.

By (1.31), (3.7), and Lemma 16, we have

**Lemma 17.** *Let* $\chi' \in \mathcal{X}_\Pi^{\mathrm{fin}}$ *and* $\Lambda' \in \mathrm{Ch}(U^0(\chi'))$ *be such that* $(\chi', \Lambda') \equiv (\chi, \Lambda)$. *Then* $\dim L_{\chi'}(\Lambda') < \infty$ *if and only if* $\dim L_\chi(\Lambda) < \infty$.

# 4 Main theorem

In this section, we keep the notation in the statement of Theorem 12. and, for $b \in J_{1,10}$, we fix $\chi_{(b)} \in \mathcal{X}_\Pi^{(b)}$, and let

$$\begin{aligned} S_{(b)} &:= \{\, \lambda \in (\mathbb{C}^\times)^N \,|\, \exists \Lambda \in \mathrm{Ch}(U^0(\chi_{(b)})) \cap \kappa_{\chi_{(b)}}^{-1}(\{\lambda\}),\ \dim L_{\chi_{(b)}}(\Lambda) < \infty \,\} \\ &= \{\, \lambda \in (\mathbb{C}^\times)^N \,|\, \forall \Lambda \in \mathrm{Ch}(U^0(\chi_{(b)})) \cap \kappa_{\chi_{(b)}}^{-1}(\{\lambda\}),\ \dim L_{\chi_{(b)}}(\Lambda) < \infty \,\}, \end{aligned} \tag{4.1}$$

where the second equality follows from Lemma 17. By Lemma 17, $S_{(b)}$ is independent from choice of $\chi' \in \mathcal{X}_\Pi^{(b)}$ with $\chi' \equiv \chi_{(b)}$.

In the statement and proof of Theorem 18 below, for $\lambda \in (\mathbb{C}^\times)^N$ and $i \in I$, let $\lambda_i$ mean the $i$-th component of $\lambda$, that is, $\lambda = (\lambda_1, \ldots, \lambda_N)$.

**Theorem 18.** *Let* $b \in J_{1,10}$. *Let* $\hat{q}$, $\hat{r}$, $\hat{\zeta}$ *be as in Theorem 12 for* $\chi_{(b)}$. *Let* $I_{(b)} := \{i \in I \,|\, \chi_{(b)}(\alpha_i, \alpha_i) \in P_\infty\}$. *Let* $S'_{(b)} := \{\lambda \in (\mathbb{C}^\times)^N \,|\, \forall j \in I_{(b)}, \exists n_j \in \mathbb{Z}_{\geq 0}, \lambda_j = \chi_{(b)}(\alpha_j, \alpha_j)^{n_j}\}$. *Then we have*

$$S_{(b)} = \begin{cases} S'_{(b)} & \text{if } b \in \{1,2,4\}, \text{ or if } b = 8 \text{ and } \hat{q}\hat{r} \notin P_\infty, \\ S''_{(b)} \uplus S'''_{(b)} & \text{otherwise,} \end{cases} \tag{4.2}$$

*where* $S''_{(b)}$ *and* $S'''_{(b)}$ *are defined as follows. Define* $\hat{v}_b \in \mathbb{C}^\times$, $n_b \in \mathbb{N}$, *and the map* $g_b : (\mathbb{C}^\times)^N \to \mathbb{C}^\times$ *by the table below.*

| $b$ | $g_b(\lambda)$ | $\hat{v}_b$ | $n_b$ |
|---|---|---|---|
| 3 | $\prod_{y\in J_{N-m,N}} \lambda_y$ | $-\hat{q}^{-1}$ | $2m$ |
| 5 | $(\prod_{y\in J_{N-m,N-2}} \lambda_y)^2\lambda_{N-1}\lambda_N$ | $\hat{q}^{-4}$ | $m$ |
| 6 | $\lambda_1^2\lambda_2^3\lambda_3^2\lambda_4$ | $\hat{q}^{-6}$ | 4 |
| 7 | $\lambda_1\lambda_2^2\lambda_3$ | $-\hat{q}^{-2}$ | 6 |
| 8 | $\lambda_1\lambda_2^2\lambda_3$ | $(\hat{q}\hat{r})^{-1}$ | 2 |
| 9 | $\lambda_1\lambda_2$ | $\hat{\zeta}\hat{q}^{-1}$ | 2 |
| 10 | $\lambda_1\lambda_2^2\lambda_3^3\lambda_4$ | $-\hat{q}^{-1}$ | 3 |

*Let* $S'_{(b);z} := \{\lambda \in S'_{(b)} \,|\, g_b(\lambda) = \hat{v}_b^z\}$. *Let*

$$S'''_{(b)} := \biguplus_{z\in J_{n_b,+\infty}} S'_{(b);z}\,. \tag{4.3}$$

*Let*

$$S''_{(3)} := \biguplus_{x\in J_{0,m-1}} \{\lambda \in S'_{(3);2x} \,|\, \forall i \in J_{N-m+x+1,N},\ \lambda_i = 1\}, \tag{4.4}$$

*Let* $g'_5(\lambda) := \prod_{i\in J_{N-m,N-1}} \lambda_i$ *for* $\lambda \in (\mathbb{C}^\times)^N$. *Let*

$$\begin{aligned} S''_{(5)} \\ := \ & \Big(\biguplus_{x\in J_{0,m-2}} \{\lambda \in S'_{(5);x} \,|\, g'_5(\lambda) = \hat{q}^{-2x},\ \forall i \in J_{N-m+x+1,N},\ \lambda_i = 1\,\}\Big) \\ & \uplus\{\lambda \in S'_{(5);m-1} \,|\, g'_5(\lambda) = \hat{q}^{-2(m-1)},\ \lambda_{N-1} = \lambda_N\}. \end{aligned} \tag{4.5}$$

*Let*

$$
\begin{aligned}
&(4.6) && S''_{(6)} &&:= \{(1,1,1,1)\} \uplus \{(\hat{q}^{-6}\lambda_3^{-1}, 1, \lambda_3, 1) \in S'_{(6);2}\} \\
& && && \quad \uplus\{(\hat{q}^{-12}\lambda_3^{-1}\lambda_4^{-2}, \hat{q}^2\lambda_4, \lambda_3, \lambda_4) \in S'_{(6);3}\}, \\
&(4.7) && S''_{(7)} &&:= \{(1,1,1)\} \uplus \{(\hat{q}^{-8}\lambda_3^{-1}, 1, \lambda_3) \in S'_{(7);4}\}, \\
&(4.8) && S''_{(8)} &&:= \{(1,1,1)\} \uplus \{\lambda \in S'_{(8);1} \,|\, \lambda_2 \in \{1, \hat{q}^{-1}\lambda_1^{-1}\}\}, \\
&(4.9) && S''_{(9)} &&:= \{(1,1)\}, \\
&(4.10) && S''_{(10)} &&:= \{(1,1,1,1)\} \uplus \{(\lambda_1, 1, 1, \lambda_4) \in S'_{(10);2} \,|\, \lambda_1\lambda_4 = -\hat{q}^{-1}\} \\
& && && \quad \uplus\{\lambda \in S'_{(10);2} \,|\, \lambda_3 \in \{1, \hat{q}^{-1}\lambda_2^{-1}\}\}.
\end{aligned}
$$

*Proof.* In this proof, let $b \in J_{1,10}$, and let $\chi_{(b)}$ be as in Theorem 12 $(b)$. In $(b)$ with $b \in J_{2,5}$ below, let $\bar{\alpha}_i$ be as in Theorem 12 $(b)$. let $N'$, $\bar{s}_i$, and $\bar{w}_0$ be those in Subsection 2.4 (defined for $\chi_{(b)}$), define $\iota \in \mathrm{Hom}_{\mathbb{Z}}(\mathbb{Z}\Pi, \mathbb{Z}^{N'})$ by $\iota(\alpha_i) := \bar{\alpha}_i$ $(i \in I)$, and let $r := r_{(b)} - \delta_{b4}$ and $f := f_{(b)}$ (see Notation 5). We also use Notations 4 and 6. Let $\chi := \chi_{(b)}$, for brevity.

(1) Assume $b = 1$. Then for any $\chi' \in \mathcal{X}_\Pi^{\mathrm{fin}}$ with $\chi' \sim \chi$, we see that $\chi' \equiv \chi$, and that $c_{ij}^{\chi'} = a_{ij}$ for all $i$, $j \in I$, where $a_{ij}$ are those one in Theorem 12 (1). Hence we have (4.2) for $b = 1$ by Lemmas 10, 15, and 16.

(2) Assume $b = 2$. We see that for any $\chi' \in \mathcal{X}_\Pi^{\mathrm{fin}}$ with $\chi' \sim \chi$, $[c_{ij}^{\chi'}]_{i,j\in I} = [c_{ij}^{\chi}]_{i,j\in I}$ and it is the Cartan matrix of type $A_N$, which implies $s_i^{\chi'} = \iota^{-1}\bar{s}_i\iota$ for all $i \in I$. By (2.18) and (2.16), we have $1^\chi s_{f,r}(\Pi) = -\Pi$. Let $r' := \frac{m(m+1)}{2} + \frac{(N-1-m)(N-m)}{2}$. We can easily see

$$
(4.11) \qquad
\begin{array}{ll}
\text{(i)} & \forall k \in I \setminus \{m+1\},\ \chi(\alpha_k, \alpha_k) \in P_\infty. \\
\text{(ii)} & \forall t_1 \in J_{1,r'},\ f(t_1) \in I \setminus \{m+1\}.\ \chi_{f,t_1-1}(\alpha_{f(t_1)}, \alpha_{f(t_1)}) \in P_\infty. \\
\text{(iii)} & \forall t_2 \in J_{r'+1,r},\ \chi_{f,t_2-1}(\alpha_{f(t_2)}, \alpha_{f(t_2)}) = -1,
\end{array}
$$

see Figure 2. Hence we have (4.2) for $b = 2$ in a way similar to that in (1).

(3) Assume $b = 3$. We see that for any $\chi' \in \mathcal{X}_\Pi^{\mathrm{fin}}$ with $\chi' \sim \chi$, $[c_{ij}^{\chi'}]_{i,j\in I} = [c_{ij}^{\chi}]_{i,j\in I}$ and it is the Cartan matrix of type $B_N$. By (2.19) and (2.16), we have $1^\chi s_{f,r}(\Pi) = -\Pi$. Let $r' := m^2$. Recall

$$
(4.12) \qquad f(t) = \begin{cases}
N - m + t & \text{if } t \in J_{1,m}, \\
f(t-m) & \text{if } t \in J_{m+1,r'}, \\
t - r' & \text{if } t \in J_{r'+1,r'+N}, \\
r' + 2N - t & \text{if } t \in J_{r'+N+1,r'+N+m}, \\
f(t-N-m) & \text{if } t \in J_{r'+N+m+1,r}.
\end{cases}
$$



$\chi_{f,t_2}$, $(t_2 \in J_{0,4})$
$\chi_{f,5}$
$\chi_{f,6}$
$\chi_{f,9}$
$\chi_{f,8}$
$\chi_{f,7}$
$\chi_{f,10}$
$\hat{q}^{-2}$ $\hat{q}^{2}$ $-1$ $\hat{q}^{-2}$ $\hat{q}^{2}$ $\hat{q}^{-2}$ $\hat{q}^{2}$
$-1$ $\hat{q}^{-2}$ $-1$ $\hat{q}^{2}$ $-1$ $\hat{q}^{-2}$ $\hat{q}^{2}$
$-1$ $\hat{q}^{-2}$ $\hat{q}^{2}$ $\hat{q}^{-2}$ $-1$ $\hat{q}^{2}$ $-1$
$\hat{q}^{2}$ $\hat{q}^{-2}$ $-1$ $\hat{q}^{2}$ $-1$ $\hat{q}^{-2}$ $-1$
$-1$ $\hat{q}^{2}$ $-1$ $\hat{q}^{-2}$ $\hat{q}^{2}$ $\hat{q}^{-2}$ $-1$
$-1$ $\hat{q}^{-2}$ $\hat{q}^{2}$ $\hat{q}^{-2}$ $\hat{q}^{2}$ $\hat{q}^{-2}$ $-1$
$\hat{q}^{2}$ $\hat{q}^{-2}$ $\hat{q}^{2}$ $\hat{q}^{-2}$ $-1$ $\hat{q}^{2}$ $\hat{q}^{-2}$
$\alpha_1$ $\alpha_2$ $\alpha_3$ $\alpha_4$
2
3
4
2
1
3

Figure 2: Dynkin diagrams of $\chi_{(2)} = \chi_{f,0}$ and $\chi_{f,u}$ with $N = 4$ and $m = 1$

where $t \in J_{1,r}$. Using Lemma 9, we can easily see

$$
(4.13)\qquad
\begin{array}{ll}
\text{(i)} & \forall k \in I \setminus \{N-m\},\ \chi(\alpha_k, \alpha_k) \in P_\infty. \\
\text{(ii)} & \forall t_1 \in J_{1,r'+(N-m-1)},\ \chi_{f,t_1} \equiv \chi. \\
\text{(iii)} & \forall t_2 \in J_{r'+(N-m),r'+(N-1)},\ \chi_{f,t_2-1}(\alpha_{f(t_2)}, \alpha_{f(t_2)}) = -1, \\
 & \text{and } \forall i \in J_{f(t_2),N},\ \forall j \in J_{f(t_2)+1,N}, \\
 & \chi_{f,t_2-1}(\alpha_i, \alpha_j)\chi_{f,t_2-1}(\alpha_j, \alpha_i) = \chi(\alpha_i, \alpha_j)\chi(\alpha_j, \alpha_i), \\
 & \chi_{f,t_2-1}(\alpha_j, \alpha_j) = \chi(\alpha_j, \alpha_j). \\
\text{(iv)} & \chi_{f,t_3-1}(\alpha_{f(t_3)}, \alpha_{f(t_3)}) = -\hat{q}^{-1}, \text{ where } t_3 := r' + N. \\
\text{(v)} & \forall t_4 \in J_{r'+N,r'+N+m},\ \chi_{f,t_4} \equiv \chi_{f,2(r'+N)-t_4-1},
\end{array}
$$

see Figure 3.

Let $\Lambda \in \mathrm{Ch}(U^0(\chi))$. For $i \in I$, let $\lambda_i := \Lambda(K_{\alpha_i}L_{-\alpha_i})$, and let $\lambda_{f,t,i} := \Lambda_{f,t}(K_{\alpha_i}L_{-\alpha_i})$ for all $t \in H(\chi, \Lambda, f)$. Let $\lambda := (\lambda_i)_{i\in I} \in (\mathbb{C}^\times)^N$. By (1.27), we see that if $\dim L_\chi(\Lambda) < \infty$, then, $\lambda \in S'_{(3)}$.

Assume that $\lambda \in S'_{(3)}$. By Lemma 15 and (4.13), we have $H(\chi, \Lambda, f) \geq r' + N - 1$, and $(\chi_{f,r'+N-m-1}, \Lambda_{f,r'+N-m-1}) \equiv (\chi, \Lambda)$. Let $\chi'' := \chi_{f,r'+N-1}$ and let $\Lambda'' := \Lambda_{f,r'+N-1}$. For $k \in J_{N-m,N}$, let

$$
(4.14)\qquad \mu_k := \begin{cases} \lambda_k & \text{if } k = N - m \text{ or } \mu_{k-1} = 1, \\ \lambda_k \hat{q}^2 \mu_{k-1} & \text{otherwise.} \end{cases}
$$

By (3.4) and (iii) of (4.13), we have $\lambda_{f,t_2-1,f(t_2)} = \mu_{f(t_2)}$, and $\lambda_{f,t_2-1,f(t_2)+1} = \lambda_{f(t_2)+1}$ for $t_2 \in J_{r'+(N-m),r'+(N-1)}$, and we have $\Lambda''(K_{\alpha_N}L_{-\alpha_N}) = \mu_N$. By (1.26) and (iv) of (4.13), we have

$$
(4.15)\qquad h_{\chi'',\Lambda'',N} < \infty \iff \exists c \in \mathbb{Z}_{\geq 0},\ \mu_N = (-\hat{q}^{-1})^c.
$$

Assume $h_{\chi'',\Lambda'',N} < \infty$. By (iii), (v) of (4.13), we have $H(\chi, \Lambda, f) \geq r' + N + m$. By Lemma 15, (3.6)-(3.7), and (v) of (4.13), we can see that

$$
(4.16)\qquad \forall t_4 \in J_{r'+N,r'+N+m},\ (\chi_{f,t_4}, \Lambda_{f,t_4}) \equiv (\chi_{f,2(r'+N)-t_4-1}, \Lambda_{f,2(r'+N)-t_4-1}).
$$



Figure 3: Dynkin diagrams of $\chi_{(3)} = \chi_{f,0}$ and $\chi_{f,u}$ with $N = 4$ and $m = 2$

In particular, $(\chi_{f,r'+N+m}, \Lambda_{f,r'+N+m}) \equiv (\chi, \Lambda)$. By (4.12), we see $H(\chi, \Lambda, f) = r$. By Lemma 16, $\dim L_{\chi_{(3)}}(\Lambda) < \infty$.

Assume that

$$\lambda \in S'_{(3)} \text{ and } \exists c_1 \in \mathbb{Z}_{\geq 0},\ g_3(\lambda) = (-\hat{q}^{-1})^{c_1}. \tag{4.17}$$

By the last paragraph, we see that $\mu_N = (-\hat{q}^{-1})^{c_1 - 2k}$ for some $k \in J_{0,m}$, and that

$$\dim L_\chi(\Lambda) < \infty \quad \Leftrightarrow \quad c_1 - 2k \geq 0. \tag{4.18}$$

Assume $c_1 \in J_{0,2m-1}$. Define $l_x \in \mathbb{Z}_{\geq 0}$ $(x \in J_{N-m+1,N})$ by $\lambda_y = \hat{q}^{2l_y}$ $(y \in J_{N-m+1,N-1})$ and $\lambda_N = \hat{q}^{l_N}$. Let $l_{N-m} := c_1 + l_N + 2\sum_{x=1}^{m-1} l_{N-m+x}$. Then $\lambda_{N-m} = (-1)^{c_1}\hat{q}^{-l_{N-m}}$. Let $Z := \{\, k \in J_{N-m+1,N} \mid \mu_{k-1} \neq 1 \,\}$. Assume $c_1 \in 2\mathbb{Z}+1$. Then $Z = J_{N-m+1,N}$. Hence $k = m$. Hence $c_1 - 2k < 0$. Assume $c_1 \in 2\mathbb{Z}$. Let $c_2 := \frac{c_1}{2}$. Let

$$\begin{aligned} c_3 &:= -l_{N-m} + 2c_2 + 2\textstyle\sum_{z=0}^{c_2} l_{N-m+z} \\ &= -(l_N + 2\textstyle\sum_{z_1=c_2+1}^{m-1} l_{N-m+z_1}). \end{aligned} \tag{4.19}$$

Since $Z \supset J_{N-m+1,N-m+c_2}$, we have $k \geq c_2$ and $\mu_{N-m+c_2} = \hat{q}^{c_3}$. If $c_3 < 0$. we have $c_2 < k$. If $c_3 = 0$, we have $c_2 = k$.

By the above argument, we see that (4.2) for $b = 3$ holds.

(4) Assume $b = 4$. Let $r' := (N-1)^2$. Recall that $f(x_1) \in J_{2,N}$ $(x_1 \in J_{1,r'})$, $f(x_2) = x_2 - r'$ $(x_2 \in J_{r'+1,r'+N})$, and $f(x_3) = r' + 2N - x_3$ $(x_3 \in J_{r'+N+1,r+1})$. See also Figure 4. Note that $s_{f(k)}^{\chi_{f,k}} = \iota^{-1}\bar{s}_{f(k)}\iota$ for all $k \in J_{1,r'+N-2}$. Define $\iota' \in \mathrm{Hom}(\mathbb{Z}\Pi, \mathbb{Z}^N)$ by $\iota'(\alpha_j) := \bar{\alpha}_j$ $(j \in J_{1,N-1})$, and $\iota'(\alpha_N) := e_{N-1} + e_N$. Then $s_{f(r'+N-1)}^{\chi_{f,r'+N-1}} = \iota^{-1}\bar{s}_{N-1}\iota'$. Define $\sigma \in \mathfrak{S}(I)$ by $\sigma(i) := i$ $(i \in J_{1,N-2})$, $\sigma(N-1) := N$, and $\sigma(N) := N-1$. Define $\tau \in \mathrm{Aut}_{\mathbb{Z}}(\mathbb{Z}\Pi)$ by $\tau(\alpha_j) := \alpha_{\sigma(j)}$. Note that $\tau s_i^{\chi'}\tau = s_{\sigma(i)}^{(\chi')^\sigma}$ holds for all $\chi' \in \mathcal{X}_\Pi$ and all $i \in I$. Note $\tau = (\iota')^{-1}\bar{s}_N\iota'$. Define $f' \in \Xi_r$ by $f'(i) := f(i)$ $(i \in J_{1,r'+N-1})$, $f'(r'+N) := N$, and $f'(j) := r+1-j$ $(j \in J_{r'+N+1,r})$. Note that $(\chi_{f',j})^\sigma \equiv \chi_{f,r+r'-j}$ $(j \in J_{r'+N-1,r})$.

We can see

$$\begin{aligned} &\iota 1^\chi s_{f',r}\tau\iota^{-1} \\ &\quad = (\iota s_{f(1)}^{\chi_{f,1}}\iota^{-1})(\iota s_{f(2)}^{\chi_{f,2}}\iota^{-1})\cdots(\iota s_{f(r'+N-2)}^{\chi_{f,r'+N-2}}\iota^{-1}) \\ &\qquad \cdot(\iota s_{N-1}^{\chi_{f,r'+N-1}}(\iota')^{-1})(\iota'\tau(\iota')^{-1})(\iota' s_{N-1}^{(\chi_{f,r'+N-2})^\sigma}\iota^{-1}) \\ &\qquad \cdot(\iota s_{N-2}^{(\chi_{f,r'+N-3})^\sigma}\iota^{-1})(\iota s_{N-3}^{(\chi_{f,r'+N-4})^\sigma}\iota^{-1})\cdots(\iota s_1^{(\chi_{f,r'})^\sigma}\iota^{-1}) \\ &\quad = (\bar{s}_{f(1)}\bar{s}_{f(2)}\cdots\bar{s}_{f(r'+N-2)})\bar{s}_{N-1}\bar{s}_N\bar{s}_{N-1}(\bar{s}_{N-2}\bar{s}_{N-3}\cdots\bar{s}_1) \\ &\quad = \bar{w}_0. \end{aligned} \tag{4.20}$$



$\chi_{f,t_2}$ ($t_2 \in J_{0,9}$)
$-1$ $\hat{q}^{-2}$ $\hat{q}^2$ $\hat{q}^{-2}$ $\hat{q}^2$ $\hat{q}^{-4}$ $\hat{q}^4$
$\alpha_1$ $\alpha_2$ $\alpha_3$ $\alpha_4$
1
$\chi_{f,10}$
$-1$ $\hat{q}^2$ $-1$ $\hat{q}^{-2}$ $\hat{q}^2$ $\hat{q}^{-4}$ $\hat{q}^4$
$\alpha_1$ $\alpha_2$ $\alpha_3$ $\alpha_4$
2
$\chi_{f,11}$
$\hat{q}^2$ $\hat{q}^{-2}$ $-1$ $\hat{q}^2$ $-1$ $\hat{q}^{-4}$ $\hat{q}^4$
$\alpha_1$ $\alpha_2$ $\alpha_3$ $\alpha_4$
3
$\chi_{f,12} \equiv (\chi_{f,12})^\sigma$
$-1$
$\hat{q}^2$ $\hat{q}^{-2}$ $\hat{q}^2$ $\hat{q}^{-2}$ $\alpha_3$
$\hat{q}^4$ $\tau$
$\alpha_1$ $\alpha_2$
$\hat{q}^{-2}$ $\alpha_4$
$-1$
4
$\chi_{f',15}$
$-1$ $\hat{q}^{-2}$ $\hat{q}^2$ $\hat{q}^{-2}$ $\hat{q}^2$ $\hat{q}^{-4}$ $\hat{q}^4$
$\alpha_1$ $\alpha_2$ $\alpha_4$ $\alpha_3$
1
$\chi_{f',14}$
$-1$ $\hat{q}^2$ $-1$ $\hat{q}^{-2}$ $\hat{q}^2$ $\hat{q}^{-4}$ $\hat{q}^4$
$\alpha_1$ $\alpha_2$ $\alpha_4$ $\alpha_3$
2
$\chi_{f',13}$
$\hat{q}^2$ $\hat{q}^{-2}$ $-1$ $\hat{q}^2$ $-1$ $\hat{q}^{-4}$ $\hat{q}^4$
$\alpha_1$ $\alpha_2$ $\alpha_4$ $\alpha_3$

Figure 4: Dynkin diagrams of $\chi_{(4)} = \chi_{f,0}$ and $\chi_{f,u}$ with $N = 4$

By (2.16) and (2.20), we have $1^\chi s_{f',r}\tau(\Pi) = -\Pi$. Hence $1^\chi s_{f',r}(\Pi) = -\Pi$. In a way similar to that in (2), we can see that (4.2) for $b = 4$ holds.

(5) Assume $b = 5$. Let $r := N^2 - m$. In a way similar to that in (4), we have $1^\chi s_{f,r}(\Pi) = -\Pi$, see Figure 5. In a way similar to that in (3), we can see that (4.2) for $b = 5$ holds.

(6)-(10) Assume $b \in J_{6,10}$. Let $r$ be the integer defined as the $(b-5)$-th



$\chi_{f,u_1}$
($u_1 \in J_{0,7} \cup \{14, 15, 22\}$)
$\hat{q}^{-2}$ $\hat{q}^2$ $-1$ $\hat{q}^{-2}$ $\hat{q}^2$ $\hat{q}^{-2}$ $\hat{q}^2$ $\alpha_4$
$\alpha_1$ $\alpha_2$ $\alpha_3$
$\hat{q}^{-2}$ $\alpha_5$ $\hat{q}^2$
2
$\chi_{f,u_2}$
($u_2 \in \{8, 13, 16, 21\}$)
$-1$ $\hat{q}^{-2}$ $-1$ $\hat{q}^2$ $-1$ $\hat{q}^{-2}$ $\hat{q}^2$ $\alpha_4$
$\alpha_1$ $\alpha_2$ $\alpha_3$
$\hat{q}^{-2}$ $\alpha_5$ $\hat{q}^2$
3
$\chi_{f,u_4}$
($u_4 \in \{10, 11, 18, 19\}$)
$-1$ $\hat{q}^{-2}$ $\hat{q}^2$ $\hat{q}^{-2}$ $\hat{q}^2$ $\hat{q}^{-2}$ $-1$ $\hat{q}^4$ $\hat{q}^{-4}$
$\alpha_1$ $\alpha_2$ $\alpha_3$ $\alpha_4$ $\alpha_5$
4
$\chi_{f,u_3}$
($u_3 \in \{9, 12, 17, 20\}$)
$-1$ $\hat{q}^{-2}$ $\hat{q}^2$ $\hat{q}^{-2}$ $-1$ $\hat{q}^2$ $-1$ $\alpha_4$
$\hat{q}^{-4}$
$\alpha_1$ $\alpha_2$ $\alpha_3$
$\hat{q}^2$ $\alpha_5$ $-1$

Figure 5: Dynkin diagrams of $\chi_{(5)} = \chi_{f,0}$ and $\chi_{f,u}$ with $N = 5$ and $m = 3$


$\chi_{f,u_1}$
$(u_1 \in J_{0,9} \cup \{18\})$
$-1$ $\hat{q}^{-2}$ $\hat{q}^{2}$ $\hat{q}^{-4}$ $\hat{q}^{4}$ $\hat{q}^{-4}$ $\hat{q}^{4}$
$\alpha_1$ $\alpha_2$ $\alpha_3$ $\alpha_4$
1
$\chi_{f,u_2}$
$(u_2 \in \{10,17\})$
$-1$ $\hat{q}^{2}$ $-1$ $\hat{q}^{-4}$ $\hat{q}^{4}$ $\hat{q}^{-4}$ $\hat{q}^{4}$
$\alpha_1$ $\alpha_2$ $\alpha_3$ $\alpha_4$
2
$\chi_{f,u_3}$
$(u_3 \in \{11,16\})$
$\hat{q}^{4}$ $\hat{q}^{-4}$ $-1$ $\hat{q}^{-2}$ $\hat{q}^{2}$ $\alpha_1$ $\hat{q}^{-2}$ $\hat{q}^{4}$ $\alpha_2$ $-1$
$\alpha_4$ $\alpha_3$
3
$\chi_{f,u_5}$
$(u_5 \in J_{13,14})$
$\hat{q}^{4}$ $\hat{q}^{-4}$ $\hat{q}^{4}$ $\hat{q}^{-4}$ $-1$ $\hat{q}^{6}$ $\hat{q}^{-6}$
$\alpha_2$ $\alpha_3$ $\alpha_4$ $\alpha_1$
4
$\chi_{f,u_4}$
$(u_4 \in \{12,15\})$
$-1$ $\hat{q}^{4}$ $\hat{q}^{-4}$ $-1$ $\hat{q}^{2}$ $\alpha_1$ $\hat{q}^{-6}$ $\hat{q}^{4}$ $\alpha_4$ $-1$
$\alpha_2$ $\alpha_3$

Figure 6: Dynkin diagrams of $\chi_{(6)} = \chi_{f,0}$ and $\chi_{f,u}$

component of $(18, 13, 7, 4, 15)$. Define $f \in \Xi_r$ by

$$(4.21)\quad (f(x))_{x\in J_{1,r}} := \begin{cases} (2,3,4,2,3,4,2,3,4,1,2,3,4,1,4,3,2,1) & \text{if } b=6,\\ (2,3,2,3,2,3,1,2,3,1,3,2,1) & \text{if } b=7,\\ (1,3,2,1,3,1,2) & \text{if } b=8,\\ (2,1,2,1) & \text{if } b=9,\\ (1,2,1,4,3,4,2,1,4,3,1,2,4,2,1) & \text{if } b=10. \end{cases}$$

Using (2.12), we can directly see that $1^{\chi} s_{f,r}(\Pi) = -\Pi$. The map $f$ has a good property similar to that for (2)-(5) in the sense that

$$(4.22)\quad \begin{aligned} &(\chi_{f,x-1}(\alpha_{f(x)}, \alpha_{f(x)}))_{x\in J_{1,r}} \\ &= \begin{cases} (\hat{q}^2, \hat{q}^4, \hat{q}^4, \hat{q}^2, \hat{q}^4, \hat{q}^4, \hat{q}^2, \hat{q}^4, \hat{q}^4, \\ -1, -1, -1, -1, \hat{q}^{-6}, -1, -1, -1, -1) \text{ if } b=6, \\ (\hat{q}^2, \hat{q}^6, \hat{q}^2, \hat{q}^6, \hat{q}^2, \hat{q}^6, \\ -1, -1, -1, -\hat{q}^{-2}, -1, -1, -1) \text{ if } b=7, \\ (\hat{q}, \hat{r}, -1, -1, (\hat{q}\hat{r})^{-1}, -1, -1) \text{ if } b=8, \\ (\hat{q}, \hat{\zeta}, \hat{\zeta}\hat{q}^{-1}, \hat{\zeta}) \text{ if } b=9, \\ (\hat{q}, \hat{q}, \hat{q}, -\hat{q}^{-1}, \\ -1, -1, -1, -1, -\hat{q}^{-1}, -\hat{q}^{-1}, -1, -1, -1, -1, -1) \text{ if } b=10. \end{cases} \end{aligned}$$

Then using $f$, in a way similar to that in (5), by tedious calculation, we can see that (4.2) for $b \in J_{6,10}$ hold.

Let us explain more precisely in the case where $b = 10$. For $\nu = (\nu_1, \nu_2, \nu_3, \nu_4) \in (J_{0,1})^4$, let $S^{\nu}_{(10)}$ be the set of elements $\lambda$ of $S_{(10)}$ such that for $\Lambda \in \mathrm{Ch}(U^0(\chi_{(10)}))$ with $\kappa_{\chi_{(10)}}(\Lambda) = \lambda$, $h_{\chi_{f,3+x},\Lambda_{f,3+x},f(4+x)} = \nu_x$ hold for all $x \in J_{1,4}$.


$\chi_{f,u_1}$
$(u_1 \in J_{0,6} \cup \{13\})$
$-1$ $\hat{q}^{-2}$ $\hat{q}^2$ $\hat{q}^{-6}$ $\hat{q}^6$
$\alpha_1$ $\alpha_2$ $\alpha_3$
1
$\chi_{f,u_2}$
$(u_2 \in \{7,12\})$
$-1$ $\hat{q}^2$ $-1$ $\hat{q}^{-6}$ $\hat{q}^6$
$\alpha_1$ $\alpha_2$ $\alpha_3$
2
$\chi_{f,u_3}$
$(u_3 \in \{8,11\})$
$\hat{q}^2$ $\alpha_1$
$-1$ $\hat{q}^{-2}$ $\hat{q}^{-4}$
$\alpha_2$ $\hat{q}^6$ $\alpha_3$ $-1$
4
$\chi_{f,u_4}$
$(u_4 J_{9,10})$
$\hat{q}^6$ $\hat{q}^{-6}$ $-1$ $\hat{q}^4$ $-\hat{q}^{-2}$
$\alpha_2$ $\alpha_3$ $\alpha_1$

Figure 7: Dynkin diagrams of $\chi_{(7)} = \chi_{f,0}$ and $\chi_{f,u}$


$\chi_{f,u_1}$
$(u_1 \in J_{0,2} \cup \{7\})$
$\hat{q}$ $\hat{q}^{-1}$ $-1$ $\hat{r}^{-1}$ $\hat{r}$
$\alpha_1$ $\alpha_2$ $\alpha_3$
2
$\chi_{f,u_2}$
$(u_2 \in \{3,6\})$
$-1$ $\alpha_1$
$-1$ $\hat{q}$ $(\hat{q}\hat{r})^{-1}$
$\alpha_2$ $\hat{r}$ $\alpha_3$ $-1$
1
$\chi_{f,u_3}$
$(u_3 \in J_{4,5})$
$\hat{q}$ $\hat{q}^{-1}$ $-1$ $\hat{q}\hat{r}$ $(\hat{q}\hat{r})^{-1}$
$\alpha_2$ $\alpha_1$ $\alpha_3$

Figure 8: Dynkin diagrams of $\chi_{(8)} = \chi_{f,0}$ and $\chi_{f,u}$


$\chi_{f,u_1}$
$(u_1 \in J_{0,1} \cup \{4\})$
$\hat{\zeta}$ $\hat{q}^{-1}$ $\hat{q}$
$\alpha_1$ $\alpha_2$
1
$\chi_{f,u_2}$
$(u_2 \in J_{2,3})$
$\hat{\zeta}$ $\hat{q}\hat{\zeta}^{-1}$ $\hat{\zeta}\hat{q}^{-1}$
$\alpha_1$ $\alpha_2$

Figure 9: Dynkin diagrams of $\chi_{(9)} = \chi_{f,0}$ and $\chi_{f,u}$


$\chi_{f,u_1}$
$(u_1 \in J_{0,4})$
$\hat{q}$ $\hat{q}^{-1}$ $\hat{q}$ $\hat{q}^{-1}$ $-1$ $-\hat{q}$ $-\hat{q}^{-1}$
$\alpha_1$ $\alpha_2$ $\alpha_3$ $\alpha_4$
3
$\chi_{f,u_2}$
$(u_2 \in \{5,13\})$
$-1$ $\alpha_3$
$\hat{q}$ $\hat{q}^{-1}$ $-1$ $\hat{q}$ $-\hat{q}^{-1}$
$\alpha_1$ $\alpha_2$ $-1$ $\alpha_4$ $-1$
4
$\chi_{f,u_3}$
$(u_3 \in \{6,12\})$
$\hat{q}$ $\hat{q}^{-1}$ $-1$ $-1$ $-1$ $-\hat{q}$ $-\hat{q}^{-1}$
$\alpha_1$ $\alpha_2$ $\alpha_4$ $\alpha_3$
2
2
$\chi_{f,14}$
$-1$ $\alpha_1$
$\hat{q}$ $\hat{q}^{-1}$ $-1$ $\hat{q}$ $-\hat{q}^{-1}$
$\alpha_3$ $\alpha_2$ $-1$ $\alpha_4$ $-1$
$\chi_{f,u_4}$
$(u_4 \in \{7,11\})$
$-1$ $\alpha_1$
$-\hat{q}^{-1}$ $-\hat{q}$ $-1$ $\hat{q}^{-1}$ $\hat{q}$
$\alpha_3$ $\alpha_4$ $-1$ $\alpha_2$ $-1$
1
1
$\chi_{f,15}$
$\hat{q}$ $\hat{q}^{-1}$ $\hat{q}$ $\hat{q}^{-1}$ $-1$ $-\hat{q}$ $-\hat{q}^{-1}$
$\alpha_3$ $\alpha_2$ $\alpha_1$ $\alpha_4$
$\chi_{f,u_5}$
$(u_5 \in J_{8,10})$
$\hat{q}$ $\hat{q}^{-1}$ $-1$ $-\hat{q}$ $-\hat{q}^{-1}$ $-\hat{q}$ $-\hat{q}^{-1}$
$\alpha_2$ $\alpha_1$ $\alpha_4$ $\alpha_3$

Figure 10: Dynkin diagrams of $\chi_{(10)} = \chi_{f,0}$ and $\chi_{f,u}$

Then we can directly see that

$$
\begin{aligned}
&S^{\nu}_{(10)} = \\
&\begin{cases}
\{\,(1,1,1,1)\,\} \quad \text{if } \nu = (0,0,0,0), \\
\{\,\lambda \in S'_{(10)} \,|\, \lambda_2 = \lambda_3 = 1,\ g_{10}(\lambda) = -\hat{q}^{-1}\,\} \quad \text{if } \nu = (0,1,1,0), \\
\{\,\lambda \in S'_{(10)} \,|\, \lambda_3 = 1,\ \exists c \in \mathbb{Z}_{\geq 0},\ g_{10}(\lambda) = (-\hat{q}^{-1})^{c+2}\,\} \\
\qquad\qquad\qquad \text{if } \nu = (0,1,1,1), \\
\{\,\lambda \in S'_{(10)} \,|\, \lambda_1\lambda_2\lambda_3^2\lambda_4 = \hat{q}^{-1},\ \exists c \in \mathbb{Z}_{\geq 0},\ g_{10}(\lambda) = (-\hat{q}^{-1})^{c+2}\,\} \\
\qquad\qquad\qquad \text{if } \nu = (1,1,1,0), \\
\{\,\lambda \in S'_{(10)} \,|\, \lambda_3 \neq 1,\ \lambda_1\lambda_2\lambda_3^2\lambda_4 \neq \hat{q}^{-1}, \\
\qquad\qquad \exists c \in \mathbb{Z}_{\geq 0},\ g_{10}(\lambda) = (-\hat{q}^{-1})^{c+3}\,\} \\
\qquad\qquad\qquad \text{if } \nu = (1,1,1,1). \\
\emptyset \quad \text{otherwise.}
\end{cases}
\end{aligned}
\tag{4.23}
$$

Then we can see that (4.2) for $b = 10$ holds. This completes the proof. □

*Remark* 7. (1) In the proof of Theorem 18 for $b \in J_{2,10}$, we have $r = |R^{+}(\chi_{(b)})|$, that is, if $b \in J_{2,3} \cup J_{5,10}$ (resp. $b = 4$), $1^{\chi}s_{f,r}$ (resp. $1^{\chi}s_{f',r}$) is a reduced expression of the longest element of the corresponding Weyl groupoid whose end domain is $\chi_{(b)}$.

(2) Let $\widehat{U}(\chi_{(b)})$ be the unital $\mathbb{C}$-algebra such that it contains $\widehat{U}(\chi_{(b)})$ and Laurent polynomial $\mathbb{C}$-algebra $X := \mathbb{C}[x_i^{\pm 1} | i \in I]$ as subalgebras, the linear map $U(\chi_{(b)}) \otimes X \to \widehat{U}(\chi_{(b)})$ $(z \otimes y \mapsto zy)$ is bijective, and $x_i K_\alpha x_i^{-1} = K_\alpha$, $x_i L_\alpha x_i^{-1} = L_\alpha$, $x_i E_j x_i^{-1} = \hat{q}^{\delta_{ij}} E_j$, $x_i F_j x_i^{-1} = \hat{q}^{-\delta_{ij}} F_j$, where recall $\hat{q} \in P_\infty$. Then we have a bijective map from $\{\Lambda \in \mathrm{Ch}(U^0(\chi_{(b)})) | \kappa_{\chi_{(b)}}(\Lambda) \in S_{(b)}\} \times (\mathbb{C}^{\times})^N$ to the set of equivalence classes of finite-dimensional $\widehat{U}(\chi_{(b)})$-modules sending $(\Lambda, \mu)$ to the equivalence class of irreducible highest weight $\widehat{U}(\chi_{(b)})$-modules with highest vectors $\widehat{v}$ such that $E_i \widehat{v} = 0$, $K_\alpha L_\beta \widehat{v} = \Lambda(K_\alpha L_\beta)\widehat{v}$ and $x_i \widehat{v} = \mu_i \widehat{v}$.

(3) It seems to be not easy to recover Theorem 18 for $b \in J_{1,8}$ from Geer's result [5] since definitions of his and our quantum groups are not so close.

(4) See [14] for deeper results concerning Theorem 18 for $b = 1$.

# References

[1] I. Angiono, On Nichols algebras of diagonal type, arXiv:1104.0268.

[2] N. Andruskiewitsch, I. Angiono and H. Yamane, On pointed Hopf superalgebras. Accepted in Contemp. Math. arXiv:1009.5148.

[3] N. Andruskiewitsch and H.-J. Schneider, Lifting of quantum linear spaces and pointed Hopf algebras of order $p^3$, J. Algebra 209 (1998) 658?691.

[4] ———, On the classification of finite-dimensional pointed Hopf algebras, Annals of Mathematics Vol. 171 (2010), No. 1, 375-417.

[5] N. Geer, Some remarks on quantized Lie superalgebras of classical type, J. Algebra 314 (2007) no. 2, 565-580

[6] I. Heckenberger, The Weyl groupoid of a Nichols algebra of diagonal type, Invent. Math. 164 (2006), 175-188.

[7] ———, Classification of arithmetic root systems, Adv. Math. 220 (2009), 59-124.

[8] ———, Lusztig isomorphisms for Drinfel'd doubles of bosonizations of Nichols algebras of diagonal type, J.Algebra, 323 (2010), 2130-2182

[9] I. Heckenberger and H. Yamane, A generalization of Coxeter groups, root systems, and Matsumoto's theorem. Math. Z. 259 (2008), 255-276.

[10] ———, Drinfel'd doubles and Shapovalov determinants, Revista de la Union Matematica Argentina vol. 51-2 (2010), pp. 107-146.

[11] V.G. Kac, Lie superalgerbras, Adv. in Math., 26 (1977) 8-96

[12] V. Kharchenko, A quantum analogue of the Poincaré-Birkhoff-Witt theorem, Algebra and Logic 38 (1999), no. 4, 259-276.

[13] G. Lusztig, *Introduction to quantum groups*, Birkhäuser, Boston, MA, 1993.

[14] D. Radford, H.-J. Schneider, On the simple representations of generalized quantum groups and quantum doubles, J. Algebra 319 (2008), 3689-3731

[15] H. Yamane, A Poincaré-Birkhoff-Witt Theorem for Quantized Universal Enveloping Algebras of Type $A_N$, Publ. RIMS, Kyoto Univ., 25 (1989), 503-520

[16] ———, Quantized enveloping algebras associated to simple Lie superalgebras and their universal R-matrices, Publ. RIMS Kyoto Univ. 30 (1994), 15-87.

[17] ———, On defining relations of affine Lie superalgebras and affine quantized universal enveloping superalgebras, Publ. RIMS Kyoto Univ. 35 (1999), no. 3, 321-390.

[18] ———, Errata to "On defining relations of affine Lie superalgebras and affine quantized universal enveloping superalgebras", Publ. RIMS Kyoto Univ. 37 (2001), no. 4, 615-619.

[19] ———, Representations of a $\mathbb{Z}/3\mathbb{Z}$-Quantum Group, Publ. RIMS, Kyoto Univ., 43 (2007), 75-93

[20] ———, Examples of the Defining Relations of the Quantum Affine Superalgebras. Available at http://www16.tok2.com/home/hiroyukipersonal/

Saeid Azam, Department of Mathematics University of Isfahan, Isfahan, Iran, P.O.Box 81745-163, azam@sci.ui.ac.ir, saeidazam@yahoo.com.

Hiroyuki Yamane, Department of Pure and Applied Mathematics, Graduate School of Information Science and Technology, Osaka University, Toyonaka, Osaka, 560-0043, Japan, yamane@ist.osaka-u.ac.jp

Malihe Yousofzadeh, Department of Mathematics University of Isfahan, Isfahan, Iran, P.O.Box 81745-163, ma.yousofzadeh@sci.ui.ac.ir.